

QUANTUM DIFFERENTIABILITY ON QUANTUM TORI

EDWARD MCDONALD, FEDOR SUKOCHEV, AND XIAO XIONG

ABSTRACT. We provide a full characterisation of quantum differentiability (in the sense of Connes) on quantum tori. We also prove a quantum integration formula which differs substantially from the commutative case.

1. INTRODUCTION

Quantum tori (also known as noncommutative tori and irrational rotation algebras) are landmark examples in noncommutative geometry. These algebras have featured in many directions in physics, such as the study of the quantum Hall effect [2, 3, 58], Matrix theory [11], string theory [52] and deformation quantisation [49]. Quantum tori have been heavily studied from the perspective of operator algebras [21, 43, 48] and were later taken as a fundamental example in noncommutative geometry (see [6], [24, Chapter 12] and [13]). In the context of foliation theory, quantum tori are studied as the C^* -algebra associated to a Kronecker foliation [9, Chapter 2, Section 9.β].

A. Connes introduced the quantised calculus in [7] as an analogue of the algebra of differential forms in a noncommutative setting, and later explored the link with the action functional of Yang-Mills theory [8]. Connes successfully applied quantised calculus in computing the Hausdorff measure of Julia sets and limit sets of Quasi-Fuchsian groups in the plane [9, Chapter 4, Section 3.γ] (for a more recent exposition see [14, 12]).

The core ingredients of the quantised calculus, as outlined in [7], are a separable Hilbert space H , a unitary self-adjoint operator F on H and a C^* -algebra \mathcal{A} represented on H such that for all $a \in \mathcal{A}$ the commutator $[F, a]$ is a compact operator on H . Then the quantised differential of $a \in \mathcal{A}$ is defined to be the operator $\bar{d}a = i[F, a]$. The compact operators on H are described by Connes as being analogous to infinitesimals, and the rate of decay of the sequence of singular values:

$$\mu(n, T) := \inf\{\|T - R\| : \text{rank}(R) \leq n\}$$

corresponds in some way to the “size” of the infinitesimal T (see [10]). In this setting one can quantify the smoothness of an element $a \in \mathcal{A}$ in terms of the rate of decay of $\{\mu(n, \bar{d}a)\}_{n=0}^{\infty}$. Of particular interest are those elements $a \in \mathcal{A}$ which satisfy:

$$\begin{aligned} \mu(n, \bar{d}a) &= O((n+1)^{-1/p}), \quad n \rightarrow \infty, \text{ or,} \\ \sum_{n=0}^{\infty} \mu(n, \bar{d}a)^p &< \infty, \text{ or,} \\ \sup_{n \geq 1} \frac{1}{\log(n+2)} \sum_{k=0}^n \mu(k, \bar{d}a)^p &< \infty, \end{aligned}$$

for some $p \in (0, \infty)$. The first condition stated above is that $\bar{d}a$ is in the weak-Schatten ideal $\mathcal{L}_{p, \infty}$, the second condition is for $\bar{d}a$ to be in the Schatten ideal \mathcal{L}_p , and the final condition is that $|\bar{d}a|^p$ is in the Macaev-Dixmier ideal $\mathcal{M}_{1, \infty}$ [9, Chapter 4, Section 2.β] (see also [37, Example 2.6.10]).

The link between quantised calculus and geometry is discussed by Connes in [8]. A model example for quantised calculus is to take a compact Riemannian spin manifold M with Dirac operator D , and define H to be the Hilbert space of square integrable sections of the spinor bundle. The algebra $\mathcal{A} = C(M)$ of continuous functions on M acts by pointwise multiplication on H , and one defines

$$F := \chi_{[0, \infty)}(D) - \chi_{(-\infty, 0)}(D).$$

2000 Mathematics Subject Classification: Primary: 46G05. Secondary: 47L10, 58B34.

Key words: Quantum tori, quantized derivative, trace formula, Sobolev space.

One then has $\bar{d}f = i[F, M_f]$, where M_f is the operator on H of pointwise multiplication by f . In quantised calculus the immediate question is to determine the relationship between the degree of differentiability of $f \in C(M)$ and the rate of decay of the singular values of $\bar{d}f$. In general, we have the following:

$$f \in C^\infty(M) \Rightarrow |\bar{d}f|^d \in \mathcal{M}_{1,\infty},$$

where d is the dimension of the manifold M [8, Theorem 3.1].

For certain special cases it is possible to obtain a far more precise understanding of the relationship between the smoothness of f and the singular values of $\bar{d}f$. The simplest example is to take the unit circle $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$, with $\mathcal{A} = C(\mathbb{T})$, $H = L_2(\mathbb{T})$ and the standard choice of F in this setting is the Hilbert transform. Then by a result of V. Peller [42, Theorem 7.3], we have that for any $p \in (0, \infty)$: $\bar{d}f \in \mathcal{L}_p$ if and only if f is in the Besov space $B_{p,p}^{1/p}(\mathbb{T})$. Peller's work has been extended to obtain even more precise relationships between f and the singular values of $\bar{d}f$, for example L. Gheorghie [22] found necessary and sufficient conditions on f to ensure that $\bar{d}f$ is in an arbitrary Riesz-Fisher space. For more details from a quantised calculus perspective, see [9, Chapter 4, Section 3.α].

In higher dimensions, the relationship between f and $\bar{d}f$ has also been studied [29, 51, 15]. To illustrate the situation, consider the d -dimensional torus \mathbb{T}^d , $d \geq 2$. The appropriate Dirac operator in this setting is:

$$D = \sum_{j=1}^d -i\gamma_j \otimes \partial_j,$$

where ∂_j denotes differentiation with respect to the j th coordinate on \mathbb{T}^d , and $\{\gamma_1, \dots, \gamma_d\}$ denotes the d -dimensional Euclidean gamma matrices, which are self-adjoint $2^{\lfloor \frac{d}{2} \rfloor} \times 2^{\lfloor \frac{d}{2} \rfloor}$ complex matrices satisfying $\gamma_j \gamma_k + \gamma_k \gamma_j = 2\delta_{j,k}1$. The operator D may be considered as an unbounded self-adjoint operator on the Hilbert space $L_2(\mathbb{T}^d, \mathbb{C}^{2^{\lfloor \frac{d}{2} \rfloor}})$. The corresponding operator F is a linear combination of Riesz transforms. The commutators of Riesz transforms and multiplication operators are studied in classical harmonic analysis: S. Janson and T. Wolff [29] proved that for $\bar{d}f$ to be in \mathcal{L}_p when $p > d$ it is necessary and sufficient that f is in the Besov space $B_{p,p}^{\frac{d}{p}}(\mathbb{T}^d)$. On the other hand, Janson and Wolff also proved that if $p \leq d$ then $\bar{d}f \in \mathcal{L}_p$ if and only if f is a constant.

A far more general characterisation of the spectral properties of commutators of Riesz transforms and multiplication operators was obtained by R. Rochberg and S. Semmes [51]. To date, investigations on the relationship between f and $\bar{d}f$ have been limited to the commutative case. To the best of our knowledge, the results treated in this paper are the first concerning quantum differentiability in the strictly noncommutative setting.

A related direction of research concerning quantised differentials is trace formulae. As early as [8] it was known that for functions on compact manifolds, it is possible to express the Dixmier trace $\text{tr}_\omega(|\bar{d}f|^p)$ as an integral of a derivative of f (See Subsection 2.1 for the relevant definitions, and [37, Chapter 6] for details on Dixmier traces).

If $f \in C^\infty(\mathbb{T}^d)$, let $\nabla f = (\partial_1 f, \partial_2 f, \dots, \partial_d f)$ be the gradient vector of f , and let $\|\nabla f\|_2 = \left(\sum_{j=1}^d |\partial_j f|^2\right)^{\frac{1}{2}}$. Then as a special case of [8, Theorem 3.3] we have:

$$(1.1) \quad \text{tr}_\omega(|\bar{d}f|^d) = k_d \int_{\mathbb{T}^d} \|\nabla f(t)\|_2^d dm(t),$$

where k_d is a constant, and m denotes the flat measure on \mathbb{T}^d (i.e., the Haar measure). From the perspective of noncommutative geometry this formula “shows how to pass from quantized 1-forms to ordinary forms, not by a classical limit, but by a direct application of the Dixmier trace” [8, Page 676]. It is also possible to prove a similar formula for functions on the non-compact manifold \mathbb{R}^d , and indeed to extend the class of traces on the left hand side of (1.1) to the much larger class of all continuous normalised traces on $\mathcal{L}_{1,\infty}$ [35].

Recently there has been work on generalising the methods of harmonic analysis on tori to quantum tori.

On a noncommutative torus \mathbb{T}_θ^d (defined in terms of an arbitrary antisymmetric real $d \times d$ matrix θ), it is possible to define analogues of many of the tools of harmonic analysis, such as differential operators and function spaces [60] (see Section 2.2). In this setting, there are analogues of all of

the components of (1.1), although the integral on the right must be replaced with the canonical trace associated to \mathbb{T}_θ^d . However the most straightforward generalisation of (1.1) to \mathbb{T}_θ^d is actually false. In this paper we state and prove a correct version of (1.1) for noncommutative tori (Theorem 1.2). The formula is stated for an appropriate class of elements $x \in L_2(\mathbb{T}_\theta^d)$ as:

$$(1.2) \quad \varphi(|\dot{d}x|^d) = c_d \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} \right) ds.$$

Here, τ is the canonical trace associated to the noncommutative torus, and c_d is a certain constant depending on d (different to the constant k_d in (1.1)). The integral is over $s = (s_1, \dots, s_d)$ in the $(d-1)$ -dimensional sphere \mathbb{S}^{d-1} , with respect to its rotation-invariant measure ds . The partial derivatives $\{\partial_1 x, \dots, \partial_d x\}$ are defined in Subsection 2.2.2. In the commutative case, the above formula reduces to (1.1) (for a full comparison, see the discussion in Subsection 1.2). There are a number of nontrivial corollaries to (1.2), which we describe in the section below.

1.1. Main results. We have three main results. We take θ to be an arbitrary $d \times d$ antisymmetric real matrix where $d \geq 2$, in particular $\theta = 0$ is not excluded. For further explanation of the notation, see Section 2 below.

Our first main result provides sufficient conditions for $\dot{d}x \in \mathcal{L}_{d,\infty}$:

Theorem 1.1. *If $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$, then $\dot{d}x$ has bounded extension, and the extension is in $\mathcal{L}_{d,\infty}$.*

The space $\dot{H}_d^1(\mathbb{T}_\theta^d)$ is a noncommutative homogeneous Sobolev space defined with respect to the partial derivatives ∂_j , $j = 1, \dots, d$ (these notions will be defined and discussed in Subsection 2.2.2). We note that the above condition is similar to that in [35, Theorem 11].

With Theorem 1.1, we can prove our second main result, the following trace formula:

Theorem 1.2. *Let $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$ be self-adjoint. Then there is a constant c_d depending only on the dimension d such that for any continuous normalised trace φ on $\mathcal{L}_{1,\infty}$ we have:*

$$\varphi(|\dot{d}x|^d) = c_d \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} \right) ds.$$

Here, the integral over \mathbb{S}^{d-1} is taken with respect to the rotation-invariant measure ds on \mathbb{S}^{d-1} , and $s = (s_1, \dots, s_d)$.

As an aside we note that it is possible to give a short argument that the integrand above is continuous as a function of $s \in \mathbb{S}^{d-1}$.

Theorem 1.2, in addition to being of interest in its own right, has a couple of corollaries, which to the best of our knowledge are novel.

Corollary 1.3. *Let $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$ be self-adjoint. Then there are constants c_d and C_d depending only on d such that for any continuous normalised trace φ on $\mathcal{L}_{1,\infty}$ we have*

$$c_d \|x\|_{\dot{H}_d^1}^d \leq \varphi(|\dot{d}x|^d) \leq C_d \|x\|_{\dot{H}_d^1}^d.$$

As a converse to Theorem 1.1, we prove our third main result: the necessity of the condition $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$ for $\dot{d}x \in \mathcal{L}_{d,\infty}$.

Theorem 1.4. *Let $x \in L_2(\mathbb{T}_\theta^d)$. If $\dot{d}x$ has bounded extension in $\mathcal{L}_{d,\infty}$ then $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$.*

The *a priori* assumption that $x \in L_2(\mathbb{T}_\theta^d)$ can be justified as follows: $L_2(\mathbb{T}_\theta^d)$ is the smallest class of x where we can define $\dot{d}x$ in a natural way. Furthermore, one can motivate this assumption by noting that an L_2 -condition is necessary and sufficient for Connes' trace theorem to hold in the commutative setting, see [36, Theorem 2.5] for details.

Since φ vanishes on the trace class \mathcal{L}_1 , Corollary 1.3 immediately yields the following noncommutative version of the $p \leq d$ component of [29, Theorem 1]:

Corollary 1.5. *If $x \in L_2(\mathbb{T}_\theta^d)$ and $\dot{d}x \in \mathcal{L}_p$, for $p \leq d$, then x is a constant.*

Indeed, the $p \leq d$ component of [29, Theorem 1] is an immediate and simple consequence of Corollary 1.5 when $\theta = 0$.

A further corollary of Theorem 1.2 is that $\varphi(|\tilde{d}x|^d)$ does not depend on the choice of continuous normalised trace φ . This implies certain asymptotic properties of the singular numbers of $\tilde{d}x$, beyond being merely in $\mathcal{L}_{d,\infty}$ [30, 53].

1.2. Comparison to the commutative case. Take $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. Consider the right hand side of the trace formula in Theorem 1.2,

$$c_d \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} \right) ds.$$

Define $\nabla x = (\partial_1 x, \partial_2 x, \dots, \partial_d x)$, and

$$\|\nabla x\|_2 := \left(\sum_{j=1}^d |\partial_j x|^2 \right)^{\frac{1}{2}}.$$

In the commutative case (when $\theta = 0$), x is a scalar valued function and $\|\nabla x\|_2^d$ coincides with the integrand in (1.1). Assuming commutativity, we can define the unit vector $u = \frac{\nabla x}{\|\nabla x\|_2}$ and take out a factor of $\|\nabla x\|_2^{\frac{d}{2}}$ to get:

$$\tau \left(\int_{\mathbb{S}^{d-1}} \left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} ds \right) = \tau \left(\|\nabla x\|_2^{\frac{d}{2}} \int_{\mathbb{S}^{d-1}} \left(\sum_{j=1}^d |u_j - s_j \sum_{k=1}^d s_k u_k|^2 \right)^{\frac{d}{2}} ds \right)$$

(where $u = (u_1, u_2, \dots, u_d)$, and the interchange of τ and the integral is easily justified by Fubini's theorem in the commutative case). However, since the measure ds on \mathbb{S}^{d-1} is invariant under rotations, we can choose coordinates $\{e_1, \dots, e_d\}$ for \mathbb{R}^d so that $u = e_1$, and then:

$$b_d := \int_{\mathbb{S}^{d-1}} \left(\sum_{j=1}^d |u_j - s_j \sum_{k=1}^d s_k u_k|^2 \right)^{\frac{d}{2}} ds$$

is independent of u , and is a constant scalar. Thus in the commutative case we have:

$$\varphi(|\tilde{d}x|^d) = c_d b_d \tau(\|\nabla x\|_2^d).$$

This recovers (1.1) upon taking $k_d = c_d b_d$. In the noncommutative case, we cannot take out a factor of $\|\nabla x\|_2^{\frac{d}{2}}$, and this explains why the form of the right hand side of Theorem 1.2 is more complicated than $k_d \tau(\|\nabla x\|_2^{\frac{d}{2}})$.

2. NOTATION

2.1. Operators, Ideals and traces. The following material concerning operator ideals and traces is standard. For more details we refer the reader to [37, 54]. Let H be a complex separable Hilbert space, and let $\mathcal{B}(H)$ denote the set of bounded operators on H , and let $\mathcal{K}(H)$ denote the ideal of compact operators on H . Given $T \in \mathcal{K}(H)$, the sequence of singular values $\mu(T) = \{\mu(k, T)\}_{k=0}^\infty$ is defined as:

$$\mu(k, T) = \inf\{\|T - R\| : \text{rank}(R) \leq k\}.$$

Equivalently, $\mu(T)$ is the sequence of eigenvalues of $|T|$ arranged in non-increasing order with multiplicities.

Let $p \in (0, \infty)$. The Schatten class \mathcal{L}_p is the set of operators T in $\mathcal{K}(H)$ such that $\mu(T)$ is p -summable, i.e. in the sequence space ℓ_p . If $p \geq 1$ then the \mathcal{L}_p norm is defined as:

$$\|T\|_p := \|\mu(T)\|_{\ell_p} = \left(\sum_{k=0}^\infty \mu(k, T)^p \right)^{1/p}.$$

With this norm \mathcal{L}_p is a Banach space, and an ideal of $\mathcal{B}(H)$.

Analogously, the weak Schatten class $\mathcal{L}_{p,\infty}$ is the set of operators T such that $\mu(T)$ is in the weak L_p -space $\ell_{p,\infty}$, with quasi-norm:

$$\|T\|_{p,\infty} = \sup_{k \geq 0} (k+1)^{1/p} \mu(k, T) < \infty.$$

As with the \mathcal{L}_p spaces, $\mathcal{L}_{p,\infty}$ is an ideal of $\mathcal{B}(H)$. We also have the following form of Hölder's inequality,

$$(2.1) \quad \|TS\|_{r,\infty} \leq c_{p,q} \|T\|_{p,\infty} \|S\|_{q,\infty}$$

where $\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$, for some constant $c_{p,q}$.

Of particular interest is $\mathcal{L}_{1,\infty}$, and we are concerned with traces on this ideal. For more details, see [37, Section 5.7] and [53]. A functional $\varphi : \mathcal{L}_{1,\infty} \rightarrow \mathbb{C}$ is called a trace if it is unitarily invariant. That is, for all unitary operators U and $T \in \mathcal{L}_{1,\infty}$ we have that $\varphi(U^*TU) = \varphi(T)$. It can then be shown that for all bounded operators B we have $\varphi(BT) = \varphi(TB)$.

An important fact about traces is that any trace φ on $\mathcal{L}_{1,\infty}$ vanishes on \mathcal{L}_1 [37, Theorem 5.7.8]. A trace φ is called continuous if it is continuous with respect to the $\mathcal{L}_{1,\infty}$ quasi-norm. It is known that not all traces on $\mathcal{L}_{1,\infty}$ are continuous [38, Remark 3.1(3)]. Within the class of continuous traces on $\mathcal{L}_{1,\infty}$ there are the well-known Dixmier traces [37, Chapter 6].

Finally, we say that a trace φ on $\mathcal{L}_{1,\infty}$ is normalised if φ takes the value 1 on any compact positive operator with eigenvalue sequence $\{\frac{1}{n+1}\}_{n=0}^\infty$ (any two such operators are unitarily equivalent, and so the particular choice of operator is inessential).

2.2. Noncommutative Tori. Harmonic analysis on noncommutative tori is an established subject. The exposition here closely follows [60], and for sake of brevity we refer the reader to [60] for a detailed exposition of the topic and provide here only the definitions relevant to this text.

2.2.1. Basic definitions. We fix an integer $d > 1$ and $\theta = \{\theta_{j,k}\}_{j,k=1}^d$, a $d \times d$ antisymmetric real matrix. The C^* -algebra of continuous functions on the noncommutative torus, denoted $C(\mathbb{T}_\theta^d)$, is the universal C^* -algebra on d unitary generators U_1, \dots, U_d which satisfy:

$$U_j U_k = e^{2\pi i \theta_{j,k}} U_k U_j, \quad 1 \leq j, k \leq d.$$

Given $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$, we adopt the shorthand notation:

$$U^n := U_1^{n_1} U_2^{n_2} \dots U_d^{n_d}.$$

There exists an action α of the torus group \mathbb{T}^d on $C(\mathbb{T}_\theta^d)$, given on a generator U_j by:

$$(2.2) \quad \alpha_z(U_j) = z_j U_j, \quad z = (z_1, z_2, \dots, z_d) \in \mathbb{T}^d.$$

The action α can be extended to a norm-continuous group of automorphisms of $C(\mathbb{T}_\theta^d)$. There is a distinguished trace state τ on $C(\mathbb{T}_\theta^d)$, which may be constructed in several ways, one of which is by averaging over α as follows: It can be shown that the fixed point subalgebra of $C(\mathbb{T}_\theta^d)$ under the action of α is exactly the trivial subalgebra $\mathbb{C}1$. Hence if $x \in C(\mathbb{T}_\theta^d)$ then averaging over \mathbb{T}^d with respect to the Haar measure m on \mathbb{T}^d :

$$\int_{\mathbb{T}^d} \alpha_z(x) dm(z)$$

yields a multiple of the identity element. Defining

$$\tau(x)1 = \int_{\mathbb{T}^d} \alpha_z(x) dm(z)$$

yields the canonical trace state τ on $C(\mathbb{T}_\theta^d)$. Given τ we can now define the GNS Hilbert space $L_2(C(\mathbb{T}_\theta^d), \tau)$, which we denote $L_2(\mathbb{T}_\theta^d)$, and we identify $C(\mathbb{T}_\theta^d)$ as an algebra of bounded operators on $L_2(\mathbb{T}_\theta^d)$, where $x \in C(\mathbb{T}_\theta^d)$ acts on $\xi \in L_2(\mathbb{T}_\theta^d)$ by left multiplication. Taking the weak operator topology closure $C(\mathbb{T}_\theta^d)''$ in $\mathcal{B}(L_2(\mathbb{T}_\theta^d))$ yields a von Neumann algebra, which we denote $L_\infty(\mathbb{T}_\theta^d)$.

The L_p -spaces for $p \in [1, \infty)$ on \mathbb{T}_θ^d are then defined as the operator L_p -spaces [44, 37] on $(L_\infty(\mathbb{T}_\theta^d), \tau)$,

$$L_p(\mathbb{T}_\theta^d) := \mathcal{L}_p(L_\infty(\mathbb{T}_\theta^d), \tau).$$

For $x \in L_1(\mathbb{T}_\theta^d)$ and $n \in \mathbb{Z}^d$, we define:

$$\widehat{x}(n) = \tau(x(U^n)^*).$$

By the definition of τ , we see that $\tau(U_n) = \delta_{n,0}$, and then standard Hilbert space arguments show that any $x \in L_2(\mathbb{T}_\theta^d)$ can be written as an L_2 -convergent series:

$$x = \sum_{n \in \mathbb{Z}^d} \widehat{x}(n) U^n,$$

with

$$(2.3) \quad \|x\|_2^2 = \sum_{n \in \mathbb{Z}^d} |\widehat{x}(n)|^2.$$

The space $C^\infty(\mathbb{T}_\theta^d)$ is defined to be the subset of $x \in C(\mathbb{T}_\theta^d)$ such that the sequence of Fourier coefficients $\{\widehat{x}(n)\}_{n \in \mathbb{Z}^d}$ has rapid decay (i.e., the sequence $\{|\widehat{x}(n)|\}_{n \in \mathbb{Z}^d}$ is eventually dominated by the reciprocal of any polynomial). We may consider $C^\infty(\mathbb{T}_\theta^d)$ as the space of smooth functions on \mathbb{T}_θ^d , since in the commutative setting this space corresponds with the space of C^∞ functions. There is also a canonical Fréchet topology on $C^\infty(\mathbb{T}_\theta^d)$, and the space $\mathcal{D}'(\mathbb{T}_\theta^d)$, called the space of distributions on \mathbb{T}_θ^d , is defined to be the topological dual of $C^\infty(\mathbb{T}_\theta^d)$.

2.2.2. *Calculus for quantum tori.* Many aspects of harmonic analysis on \mathbb{T}^d carry over to \mathbb{T}_θ^d . For example we may define the partial differentiation operators ∂_j , $j = 1, \dots, d$ by:

$$\partial_j(U^n) = 2\pi i n_j U^n, \quad n = (n_1, \dots, n_d) \in \mathbb{Z}^d.$$

Every partial derivation ∂_j can be viewed a densely defined closed (unbounded) operator on $L_2(\mathbb{T}_\theta^d)$, whose adjoint is equal to $-\partial_j$. Let $\Delta = \partial_1^2 + \dots + \partial_d^2$ be the Laplacian. Then $\Delta = -(\partial_1^* \partial_1 + \dots + \partial_d^* \partial_d)$, so $-\Delta$ is a positive operator on $L_2(\mathbb{T}_\theta^d)$ with spectrum equal to $\{4\pi^2 |n|^2 : n \in \mathbb{Z}^d\}$. As in the Euclidean case, we let $D_j = -i\partial_j$, which is then self-adjoint. Given $n = (n_1, \dots, n_d) \in \mathbb{N}_0^d$ (\mathbb{N}_0 denoting the set of nonnegative integers), the associated partial derivation D^n is defined to be $D_1^{n_1} \dots D_d^{n_d}$. The order of D^n is $|n|_1 = n_1 + \dots + n_d$. By duality, the derivations transfer to $\mathcal{D}'(\mathbb{T}_\theta^d)$ as well.

For $\alpha \in \mathbb{R}$, denote by J^α the α -order Bessel potential $(1 - \Delta)^{\frac{\alpha}{2}}$. The potential (or fractional) Sobolev space of order $\alpha \in \mathbb{R}$ is defined to be

$$(2.4) \quad H_p^\alpha(\mathbb{T}_\theta^d) = \{x \in \mathcal{D}'(\mathbb{T}_\theta^d) : J^\alpha x \in L_p(\mathbb{T}_\theta^d)\},$$

equipped with the norm

$$\|x\|_{H_p^\alpha} = \|J^\alpha x\|_p.$$

Since J^0 is the identity, $H_p^0(\mathbb{T}_\theta^d) = L_p(\mathbb{T}_\theta^d)$. As in the classical case, if α is a non-negative integer then $H_p^\alpha(\mathbb{T}_\theta^d)$ admits an equivalent norm in terms of the sum of the p -norms of the partial derivatives of order up to α . To be explicit, the Sobolev space of order $k \in \mathbb{N}$ on \mathbb{T}_θ^d may be described as:

$$H_p^k(\mathbb{T}_\theta^d) = \{x \in \mathcal{D}'(\mathbb{T}_\theta^d) : D^n x \in L_p(\mathbb{T}_\theta^d) \text{ for each } n \in \mathbb{N}_0^d \text{ with } |n|_1 \leq k\},$$

equipped with the norm

$$\|x\|_{H_p^k} = \left(\sum_{0 \leq |n|_1 \leq k} \|D^n x\|_p^p \right)^{\frac{1}{p}}.$$

The equivalence of the above norm and the Bessel potential norm $\|J^k x\|_p$ is a well-established fact in the theory of harmonic analysis on \mathbb{T}_θ^d , being proved in the $p = 2$ case by [55, Theorem 2.1] and a later proof for general p can be found as [60, Theorem 2.9].

In this paper, we will mainly use the ‘‘homogeneous’’ Sobolev space $\dot{H}_p^1(\mathbb{T}_\theta^d)$ and the potential Sobolev spaces $H_2^\alpha(\mathbb{T}_\theta^d)$. The norm of $\dot{H}_p^1(\mathbb{T}_\theta^d)$ with $p \geq 2$, may be described in the following equivalent forms:

$$(2.5) \quad \|x\|_{\dot{H}_p^1} = \left(\sum_{j=1}^d \|\partial_j x\|_p^p \right)^{\frac{1}{p}} \approx \sum_{j=1}^d \|\partial_j x\|_p \approx \left\| \left(\sum_{j=1}^d |\partial_j x|^2 \right)^{\frac{1}{2}} \right\|_p,$$

where the relevant constants depend only on d and p . Then $\dot{H}_p^1(\mathbb{T}_\theta^d)$ may be defined as the subspace of $\mathcal{D}'(\mathbb{T}_\theta^d)$ for which the above norm is finite. Note that the difference between $\dot{H}_p^1(\mathbb{T}_\theta^d)$ and $H_p^1(\mathbb{T}_\theta^d)$ is that for $\dot{H}_p^1(\mathbb{T}_\theta^d)$ we do not assume that the L_p -norm is finite. For any $x \in H_p^1(\mathbb{T}_\theta^d)$, we have the following Poincaré type inequality

$$(2.6) \quad \|x - \widehat{x}(0)\|_p \leq C_{p,d} \|x\|_{\dot{H}_p^1}.$$

See [60, Theorem 2.12]. For every $\alpha \in \mathbb{R}$, the space $H_2^\alpha(\mathbb{T}_\theta^d)$ is a Hilbert space with the inner product

$$\langle x, y \rangle = \tau(J^\alpha y^* J^\alpha x).$$

It is proved in [55, Theorem 3.3] and [27, Proposition 9.2] for arbitrary real $\alpha, \beta \in \mathbb{R}$ with $\alpha > \beta$, the embedding

$$(2.7) \quad H_2^\alpha(\mathbb{T}_\theta^d) \hookrightarrow H_2^\beta(\mathbb{T}_\theta^d) \quad \text{is compact.}$$

The Dirac operator D (more precisely, the spin-Dirac operator) is defined in terms of γ matrices in direct analogy to commutative tori. Define $N = 2^{\lfloor \frac{d}{2} \rfloor}$ and select $N \times N$ complex self-adjoint matrices $\{\gamma_1, \dots, \gamma_d\}$ satisfying $\gamma_j \gamma_k + \gamma_k \gamma_j = 2\delta_{j,k}1$, and define:

$$D = \sum_{j=1}^d \gamma_j \otimes D_j$$

as an unbounded, densely defined linear operator on the Hilbert space $\mathbb{C}^N \otimes L_2(\mathbb{T}_\theta^d)$. This definition coincides with [24, Definition 12.14].

Since all D_j 's are self-adjoint, D is also self-adjoint. We then define the sign of D via the Borel functional calculus, which can be expressed as

$$\text{sgn}(D) = \sum_{j=1}^d \gamma_j \otimes \frac{D_j}{\sqrt{D_1^2 + D_2^2 + \dots + D_d^2}}.$$

Given $x \in L_\infty(\mathbb{T}_\theta^d)$, denote by $M_x : y \mapsto xy$ the operator of left multiplication on $L_2(\mathbb{T}_\theta^d)$. The operator $1 \otimes M_x$ is a bounded linear operator on $\mathbb{C}^N \otimes L_2(\mathbb{T}_\theta^d)$, where 1 denotes the identity operator on \mathbb{C}^N . The commutator

$$\overline{d}x := i[\text{sgn}(D), 1 \otimes M_x], \quad x \in L_\infty(\mathbb{T}_\theta^d)$$

denotes the quantised differential on quantum tori.

On the other hand, if x is not necessarily bounded we may still define $\overline{d}x$ on the dense subspace $C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$ as follows. Suppose that $x \in L_2(\mathbb{T}_\theta^d)$. Then if $\eta \in C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$, we will have $(1 \otimes M_x)\eta \in L_2(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$. Moreover, $\text{sgn}(D)\eta$ is still in $C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$ since by definition an element of $C^\infty(\mathbb{T}_\theta^d)$ has Fourier coefficients of rapid decay, and $\text{sgn}(D)$ is represented as a Fourier multiplier with bounded symbol. Thus the expression:

$$(\overline{d}x)\eta := i\text{sgn}(D)(1 \otimes M_x)\eta - i(1 \otimes M_x)\text{sgn}(D)\eta$$

is a well-defined element of $L_2(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$ for all $\eta \in C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$.

2.2.3. Fourier multipliers for quantum tori. Let g be a bounded scalar function on \mathbb{Z}^d . For $x \in L_2(\mathbb{T}_\theta^d)$, the Fourier multiplier T_g with symbol g is defined on x by:

$$(2.8) \quad T_g x = \sum_{n \in \mathbb{Z}^d} g(n) \widehat{x}(n) U^n.$$

By virtue of the Plancherel identity (2.3), T_g indeed defines a bounded linear operator on $L_2(\mathbb{T}_\theta^d)$ and the above series converges in the L_2 -sense. If g is unbounded, we may define T_g on the dense subspace of $L_2(\mathbb{T}_\theta^d)$ of those x with finitely many non-zero Fourier coefficients.

An equivalent perspective on Fourier series is to consider a function $\phi \in L_1(\mathbb{T}^d)$ on the commutative torus. We may then define the convolution of ϕ with $x \in L_2(\mathbb{T}_\theta^d)$ by:

$$\phi * x = \int_{\mathbb{T}^d} \alpha_w(x) \phi(w) dw.$$

In terms of Fourier coefficients, we have:

$$\widehat{\phi * x} = T_{\widehat{\phi}} x.$$

Fourier multipliers for quantum tori were studied in detail in [60, Chapter 7] (there, T_g was denoted by M_g). From the perspective of functional calculus, we may also write:

$$T_g = g\left(\frac{1}{2\pi i} \partial_1, \frac{1}{2\pi i} \partial_2, \dots, \frac{1}{2\pi i} \partial_d\right).$$

The above defined derivatives D^α , Laplacian Δ , and Bessel potential J^α may all be viewed as Fourier multipliers: the symbol of D_j is $2\pi\xi_j$; the symbol of Δ is $-|2\pi\xi|^2$; and the symbol of J^α is $(1 + |2\pi\xi|^2)^{\frac{\alpha}{2}}$. We will denote by $\langle \xi \rangle$ the function $(1 + |\xi|^2)^{\frac{1}{2}}$ in the sequel.

A far reaching extension of the notion of a Fourier multiplier is a pseudodifferential operator. We outline the pseudodifferential operator theory for the noncommutative torus in Section 5.

3. CWIKEL-TYPE ESTIMATES FOR QUANTUM TORI

In the classical, commutative setting, Cwikel estimates are bounds on the singular values of operators of the form:

$$M_f g(-i\nabla)$$

where f and g are essentially bounded functions on \mathbb{R}^d , and M_f and $g(-i\nabla)$ denote pointwise multiplication and Fourier multiplication on $L_2(\mathbb{R}^d)$ respectively (see e.g. [54, Chapter 4] and [17]).

In the setting of noncommutative tori, we instead consider operators of the form $M_x T_g$, where $x \in L_\infty(\mathbb{T}_\theta^d)$ and $g \in \ell_\infty(\mathbb{Z}^d)$. We can obtain the following as a special case of [33]:

Theorem 3.1. (i) *If $x \in L_p(\mathbb{T}_\theta^d)$ and $g \in \ell_p(\mathbb{Z}^d)$ with $2 \leq p < \infty$, then $M_x T_g$ is in \mathcal{L}_p and*

$$\|M_x T_g\|_{\mathcal{L}_p} \leq C_p \|x\|_p \|g\|_p.$$

(ii) *If $x \in L_p(\mathbb{T}_\theta^d)$ and $g \in \ell_{p,\infty}(\mathbb{Z}^d)$ with $2 < p < \infty$, then $M_x T_g$ is in $\mathcal{L}_{p,\infty}$ and*

$$\|M_x T_g\|_{\mathcal{L}_{p,\infty}} \leq C_p \|x\|_p \|g\|_{p,\infty}.$$

Proof. We in fact prove the following far stronger estimate, stated in the language of symmetric function spaces [37, Chapter 2]: For any symmetric function space E whose norm satisfies the Fatou property¹ and is an interpolation space of L_2 and L_∞ , if $x \otimes g \in E(L_\infty(\mathbb{T}_\theta^d) \otimes \ell_\infty(\mathbb{Z}^d))$ then $M_x T_g$ is in $E(\mathcal{B}(L_2(\mathbb{T}_\theta^d)))$, with norm bound,

$$(3.1) \quad \|M_x T_g\|_{E(\mathcal{B}(L_2(\mathbb{T}_\theta^d)))} \leq C_E \|x \otimes g\|_{E(L_\infty(\mathbb{T}_\theta^d) \otimes \ell_\infty(\mathbb{Z}^d))}.$$

After proving (3.1), we explain how it entails the results in the statement of the theorem.

In fact (3.1) can be obtained by a direct application of [33, Corollary 3.5]. Here we have two von Neumann algebras $L_\infty(\mathbb{T}_\theta^d)$ and $\ell_\infty(\mathbb{Z}^d)$ represented on the same Hilbert space $L_2(\mathbb{T}_\theta^d)$ by left multiplication and Fourier multiplication respectively. In this setting, we can use [33, Corollary 3.5] which states that if we have an estimate of the form:

$$(3.2) \quad \|M_x T_g\|_{\mathcal{L}_2(\mathcal{B}(L_2(\mathbb{T}_\theta^d)))} \leq \|x\|_{L_2(\mathbb{T}_\theta^d)} \|g\|_{\ell_2(\mathbb{Z}^d)}$$

then (3.1) follows.

To prove (3.2), we can express the Hilbert-Schmidt norm in terms of an expansion with respect to the basis $\{U^m\}_{m \in \mathbb{Z}^d}$ of $L_2(\mathbb{T}_\theta^d)$,

$$\begin{aligned} \|M_x T_g\|_{\mathcal{L}_2}^2 &= \sum_{m,n \in \mathbb{Z}^d} \left| \tau \left(x (T_g U^m) (U^n)^* \right) \right|^2 \\ &= \sum_{m,n \in \mathbb{Z}^d} \left| \tau \left(x g(m) U^m (U^n)^* \right) \right|^2 = \sum_{m,n \in \mathbb{Z}^d} |g(m)|^2 \left| \tau \left(x U^m (U^n)^* \right) \right|^2 \\ &= \sum_{m \in \mathbb{Z}^d} |g(m)|^2 \sum_{n \in \mathbb{Z}^d} \left| \tau \left(x U^m (U^n)^* \right) \right|^2. \end{aligned}$$

By the Plancherel formula (2.3), we have

$$\sum_{n \in \mathbb{Z}^d} \left| \tau \left(x U^m (U^n)^* \right) \right|^2 = \|x U^m\|_2^2 = \|x\|_2^2.$$

Thus,

$$\|M_x T_g\|_{\mathcal{L}_2}^2 = \|x\|_2^2 \|g\|_2^2.$$

Hence, (3.2) holds and thus by [33, Corollary 3.5] it follows that (3.1) holds.

Now, we take $E = L_p$ in (3.1) for $p \in (2, \infty)$. This is indeed an interpolation space between L_2 and L_∞ whose norm satisfies the Fatou property. Then combining (3.1) with the identity

$$\|x \otimes g\|_{L_p(L_\infty(\mathbb{T}_\theta^d) \otimes \ell_\infty(\mathbb{Z}^d))} = \|x\|_p \|g\|_p$$

¹meaning that if A_n is a sequence of positive operators with $A_n \uparrow A$ in the weak operator topology, then $\|A\|_E \leq \sup_n \|A_n\|_E$

yields (i).

Finally, to obtain (ii), we take $E = L_{p,\infty}$ in (3.1) and use the estimate:

$$\|x \otimes g\|_{L_{p,\infty}(L_\infty(\mathbb{T}_\theta^d) \otimes \ell_\infty(\mathbb{Z}^d))} \leq \|x\|_p \|g\|_{p,\infty}.$$

This completes the proof of (ii). \square

Consider the function on \mathbb{Z}^d , $n \mapsto (1 + |n|^2)^{-\frac{d}{2}}$. When $|n| > 1$, we have $(1 + |n|^2)^{-\frac{d}{2}} \leq |n|^{-d}$. For $|n| \leq 1$, $(1 + |n|^2)^{-\frac{d}{2}}$ is bounded from above by 1. Hence $n \mapsto (1 + |n|^2)^{-\frac{d}{2}} \in \ell_{1,\infty}(\mathbb{Z}^d)$, and so $n \mapsto (1 + |n|^2)^{-\frac{\beta}{2}} \in \ell_{\frac{d}{\beta},\infty}(\mathbb{Z}^d)$. Then it follows immediately from the above theorem that

Corollary 3.2. *Consider the linear operator $(1 \otimes x)(1 + D^2)^{-\frac{\beta}{2}}$ on $\mathbb{C}^N \otimes L_2(\mathbb{T}_\theta^d)$. If $x \in L_{\frac{d}{\beta}}(\mathbb{T}_\theta^d)$ with $\frac{d}{\beta} > 2$, then $(1 \otimes M_x)(1 + D^2)^{-\frac{\beta}{2}} \in \mathcal{L}_{\frac{d}{\beta},\infty}$, and*

$$\|(1 \otimes M_x)(1 + D^2)^{-\frac{\beta}{2}}\|_{\mathcal{L}_{\frac{d}{\beta},\infty}} \leq C \|x\|_{\frac{d}{\beta}},$$

where the constant $C > 0$ depends only on d and β .

At this point it is worth noting that since the function $n \mapsto (1 + |n|^2)^{-\frac{\alpha}{2}}$ is in $\ell_{\frac{d}{\alpha},\infty}(\mathbb{Z})$, for all $\alpha > 0$ we have:

$$(3.3) \quad J^{-\alpha} \in \mathcal{L}_{\frac{d}{\alpha},\infty}.$$

4. PROOF OF THEOREM 1.1

This section is devoted to the proof of Theorem 1.1, that is, that the condition $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$ is sufficient for $\bar{d}x \in \mathcal{L}_{d,\infty}$, and with an explicit norm bound:

$$\|\bar{d}x\|_{d,\infty} \leq C_d \|x\|_{\dot{H}_d^1(\mathbb{T}_\theta^d)}.$$

Note that due to the Poincaré inequality (2.6), $\dot{H}_d^1(\mathbb{T}_\theta^d)$ is a subset of $L_d(\mathbb{T}_\theta^d)$, and thus in particular the operator $\bar{d}x$ is well-defined.

The following lemma is a corollary of Theorem 3.1(i).

Lemma 4.1. *Suppose that $p > \frac{d}{2}$ and $x \in L_p(\mathbb{T}_\theta^d)$. If $p \geq 2$, then there exists a constant $C_{p,d} > 0$ such that*

$$\left\| \left[\text{sgn}(D) - \frac{D}{\sqrt{1+D^2}}, 1 \otimes M_x \right] \right\|_{\mathcal{L}_p} \leq C_{p,d} \|x\|_p,$$

meaning that, if $x \in L_p(\mathbb{T}_\theta^d)$ then the above commutator (initially defined on $C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$) admits an extension to a bounded operator which is in the ideal \mathcal{L}_p with the above norm bound.

Proof. Let $1 \leq j \leq d$, and for $n \in \mathbb{Z}^d$ define

$$h_j(n) := \frac{n_j}{|n|} - \frac{n_j}{((2\pi)^{-2} + |n|^2)^{\frac{1}{2}}}.$$

Thus,

$$T_{h_j} = h_j \left(-\frac{i}{2\pi} \nabla \right) = \frac{-i\partial_j}{\sqrt{-\Delta}} - \frac{-i\partial_j}{(1-\Delta)^{\frac{1}{2}}}$$

and so,

$$\begin{aligned} \text{sgn}(D) - \frac{D}{\sqrt{1+D^2}} &= \sum_{j=1}^d \gamma_j \otimes \left(\frac{-i\partial_j}{\sqrt{-\Delta}} - \frac{-i\partial_j}{(1-\Delta)^{\frac{1}{2}}} \right) \\ &= \sum_{j=1}^d \gamma_j \otimes h_j \left(-\frac{i}{2\pi} \nabla \right) \\ &= \sum_{j=1}^d \gamma_j \otimes T_{h_j}. \end{aligned}$$

One can easily check that $h_j \in \ell_p(\mathbb{Z}^d)$ as $p > \frac{d}{2}$. Expanding out the commutator,

$$\begin{aligned} \left[\operatorname{sgn}(D) - \frac{D}{\sqrt{1+D^2}}, 1 \otimes M_x \right] &= \left[\sum_{j=1}^d \gamma_j \otimes T_{h_j}, 1 \otimes M_x \right] \\ &= \sum_{j=1}^d \gamma_j \otimes [T_{h_j}, M_x]. \end{aligned}$$

Hence,

$$\begin{aligned} \left\| \left[\operatorname{sgn}(D) - \frac{D}{\sqrt{1+D^2}}, 1 \otimes M_x \right] \right\|_{\mathcal{L}_p} &\leq d \max_{1 \leq j \leq d} \| [T_{h_j}, M_x] \|_{\mathcal{L}_p} \\ &\leq d \max_{1 \leq j \leq d} (\| T_{h_j} M_x \|_{\mathcal{L}_p} + \| M_x T_{h_j} \|_{\mathcal{L}_p}) \\ &= d \max_{1 \leq j \leq d} (\| M_x^* T_{h_j} \|_{\mathcal{L}_p} + \| M_x T_{h_j} \|_{\mathcal{L}_p}). \end{aligned}$$

The desired conclusion follows then from Theorem 3.1.(i). \square

The proof of the next lemma relies on the technique of double operator integrals (see [41] and [46] and references therein). Let H be a (complex) separable Hilbert space. Let D_0 and D_1 be self-adjoint (potentially unbounded) operators on H , and E^0 and E^1 be the associated spectral measures. For all $x, y \in \mathcal{L}_2(H)$, the measure $(\lambda, \mu) \mapsto \operatorname{Tr}(x dE^0(\lambda) y dE^1(\mu))$ is a countably additive complex valued measure on \mathbb{R}^2 . We say that $\phi \in L_\infty(\mathbb{R}^2)$ is $E^0 \otimes E^1$ integrable if there exists an operator $T_\phi^{D_0, D_1} \in \mathcal{B}(\mathcal{L}_2(H))$ such that for all $x, y \in \mathcal{L}_2(H)$,

$$\operatorname{Tr}(x T_\phi^{D_0, D_1} y) = \int_{\mathbb{R}^2} \phi(\lambda, \mu) \operatorname{Tr}(x dE^0(\lambda) y dE^1(\mu)).$$

The operator $T_\phi^{D_0, D_1}$ is called the transformer. For $A \in \mathcal{L}_2(H)$, we define

$$(4.1) \quad T_\phi^{D_0, D_1}(A) = \int_{\mathbb{R}^2} \phi(\lambda, \mu) dE^0(\lambda) A dE^1(\mu).$$

This is called a double operator integral.

Lemma 4.2. *Let $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. Then*

$$\left\| \left[\frac{D}{\sqrt{1+D^2}}, 1 \otimes M_x \right] \right\|_{\mathcal{L}_{d,\infty}} \leq B_d \|x\|_{\dot{H}_d^1}$$

where the constant $B_d > 0$ depends only on d . As with Lemma 4.1, the above commutator is interpreted as being initially defined on $C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$.

Proof. Set $g(t) = t(1+t^2)^{-\frac{1}{2}}$ for $t \in \mathbb{R}$. Suppose initially that $x \in C^\infty(\mathbb{T}_\theta^d)$. Under this assumption, $[D, 1 \otimes M_x]$ extends to a bounded operator, and thus we can apply [4, Theorem 4.1] (see also Proposition 2.6 and Theorem 3.1 in [45]) to get

$$(4.2) \quad [g(D), 1 \otimes M_x] = T_{g^{[1]}}^{D,D}([D, 1 \otimes M_x]),$$

where $g^{[1]}(\lambda, \mu) := \frac{g(\lambda) - g(\mu)}{\lambda - \mu}$ for different $\lambda, \mu \in \mathbb{R}$. By [35, Lemma 9], we have $g^{[1]} = \psi_1 \psi_2 \psi_3$, with

$$\psi_1 = 1 + \frac{1 - \lambda\mu}{(1 + \lambda^2)^{\frac{1}{2}}(1 + \mu^2)^{\frac{1}{2}}}, \quad \psi_2 = \frac{(1 + \lambda^2)^{\frac{1}{4}}(1 + \mu^2)^{\frac{1}{4}}}{(1 + \lambda^2)^{\frac{1}{2}} + (1 + \mu^2)^{\frac{1}{2}}}, \quad \psi_3 = \frac{1}{(1 + \lambda^2)^{\frac{1}{4}}(1 + \mu^2)^{\frac{1}{4}}}.$$

It follows that

$$(4.3) \quad T_{g^{[1]}}^{D,D} = T_{\psi_1}^{D,D} T_{\psi_2}^{D,D} T_{\psi_3}^{D,D}.$$

By [35, Lemma 8], we see that the transformer $T_{\psi_2}^{D,D}$ is bounded on both \mathcal{L}_1 and \mathcal{L}_∞ .

For $k = 1, 3$ the function ψ_k can be written as a linear combination of products of bounded functions of λ and of μ , and from this it follows that $T_{\psi_k}^{D,D}$ is a bounded linear map on \mathcal{L}_1 and \mathcal{L}_∞ . For further details, see e.g. [46, Corollary 2] and [50, Corollary 2.4].

Then by real interpolation of $(\mathcal{L}_1, \mathcal{L}_\infty)$ (see [18] or [61]), the transformers $T_{\psi_k}^{D,D}$ with $k = 1, 2, 3$ are bounded linear transformations from $\mathcal{L}_{d,\infty}$ to $\mathcal{L}_{d,\infty}$.

We now exploit the identity in (4.2) and the product of terms in (4.3), noticing that

$$\begin{aligned} \| [g(D), 1 \otimes M_x] \|_{\mathcal{L}_{d,\infty}} &\leq \| T_{\psi_1}^{D,D} \|_{\mathcal{L}_{d,\infty} \rightarrow \mathcal{L}_{d,\infty}} \| T_{\psi_2}^{D,D} \|_{\mathcal{L}_{d,\infty} \rightarrow \mathcal{L}_{d,\infty}} \\ &\quad \times \| T_{\psi_3}^{D,D} ([D, 1 \otimes M_x]) \|_{\mathcal{L}_{d,\infty}} \\ &\leq C_d \| T_{\psi_3}^{D,D} ([D, 1 \otimes M_x]) \|_{\mathcal{L}_{d,\infty}}, \end{aligned}$$

where the constant $C_d > 0$ does not depend on x . Since $\psi_3(\lambda, \mu) = (1 + \lambda^2)^{-1/4} (1 + \mu^2)^{-1/4}$ is a product a function of λ and a function of μ , by (4.1), we have

$$T_{\psi_3}^{D,D} ([D, 1 \otimes M_x]) = (1 + D^2)^{-1/4} [D, 1 \otimes M_x] (1 + D^2)^{-1/4}.$$

Hence

$$\| [g(D), 1 \otimes M_x] \|_{\mathcal{L}_{d,\infty}} \leq C_d \| (1 + D^2)^{-1/4} [D, 1 \otimes M_x] (1 + D^2)^{-1/4} \|_{\mathcal{L}_{d,\infty}}.$$

Expanding out D and using the quasi-triangle inequality for $\mathcal{L}_{d,\infty}$, we have

$$\begin{aligned} &\| (1 + D^2)^{-1/4} [D, 1 \otimes M_x] (1 + D^2)^{-1/4} \|_{\mathcal{L}_{d,\infty}} \\ &\leq K_d \sum_{j=1}^d \| (1 + D^2)^{-1/4} [\gamma_j \otimes D_j, 1 \otimes M_x] (1 + D^2)^{-1/4} \|_{\mathcal{L}_{d,\infty}}, \end{aligned}$$

where $K_d > 0$ depends only on d . But $[\gamma_j \otimes D_j, 1 \otimes M_x] = -i\gamma_j \otimes M_{\partial_j x}$, thus we obtain

$$\| (1 + D^2)^{-1/4} [\gamma_j \otimes D_j, 1 \otimes M_x] (1 + D^2)^{-1/4} \|_{\mathcal{L}_{d,\infty}} = \| (1 - \Delta)^{-1/4} M_{\partial_j x} (1 - \Delta)^{-1/4} \|_{\mathcal{L}_{d,\infty}}.$$

Note that the first norm $\| \cdot \|_{\mathcal{L}_{d,\infty}}$ is the norm of $\mathcal{L}_{d,\infty}(\mathbb{C}^N \otimes L_2(\mathbb{T}_\theta^d))$, and the second one is the norm of $\mathcal{L}_{d,\infty}(L_2(\mathbb{T}_\theta^d))$.

We are reduced to estimating the quantity $\| (1 - \Delta)^{-1/4} M_{\partial_j x} (1 - \Delta)^{-1/4} \|_{\mathcal{L}_{d,\infty}}$. By polar decomposition, for every j , there is a partial isometry U_j such that

$$\partial_j x = U_j |\partial_j x| = U_j |\partial_j x|^{\frac{1}{2}} |\partial_j x|^{\frac{1}{2}}.$$

Taking $\beta = \frac{1}{2}$, and recalling that x is such that $\| U_j |\partial_j x|^{\frac{1}{2}} \|_{2d} \leq \| |\partial_j x|^{\frac{1}{2}} \|_{2d} = \| \partial_j x \|_d^{\frac{1}{2}} < \infty$, we apply Corollary 3.2 to get (for some constant Q_d)

$$\| M_{|\partial_j x|^{\frac{1}{2}}} (1 - \Delta)^{-1/4} \|_{\mathcal{L}_{2d,\infty}} = \| (1 - \Delta)^{-1/4} M_{|\partial_j x|^{\frac{1}{2}}} \|_{\mathcal{L}_{2d,\infty}} \leq Q_d \| |\partial_j x|^{\frac{1}{2}} \|_{2d}$$

and

$$\| (1 - \Delta)^{-1/4} M_{U_j |\partial_j x|^{\frac{1}{2}}} \|_{\mathcal{L}_{2d,\infty}} \leq Q_d \| U_j |\partial_j x|^{\frac{1}{2}} \|_{2d} \leq Q_d \| |\partial_j x|^{\frac{1}{2}} \|_{2d}.$$

Thus, by the Hölder inequality (2.1),

$$\| (1 - \Delta)^{-1/4} M_{\partial_j x} (1 - \Delta)^{-1/4} \|_{\mathcal{L}_{d,\infty}} \leq c_{\frac{d}{2}, \frac{d}{2}} Q_d^2 \| |\partial_j x|^{\frac{1}{2}} \|_{2d}^2 = c_{\frac{d}{2}, \frac{d}{2}} Q_d^2 \| \partial_j x \|_d.$$

Taking $B_d = c_{\frac{d}{2}, \frac{d}{2}} d Q_d^2 C_d K_d$, we conclude that

$$(4.4) \quad \| [g(D), 1 \otimes M_x] \|_{\mathcal{L}_{d,\infty}} \leq B_d \sum_{j=1}^d \| \partial_j x \|_d \leq B_d \| x \|_{\dot{H}_d^1}.$$

We now remove the initial assumption that $x \in C^\infty(\mathbb{T}_\theta^d)$. Suppose that $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. As $C^\infty(\mathbb{T}_\theta^d)$ is dense in $H_d^1(\mathbb{T}_\theta^d)$ [60, Proposition 2.7], we may select a sequence $\{x_n\}_{n=0}^\infty \subset C^\infty(\mathbb{T}_\theta^d)$ such that $\lim_{n \rightarrow \infty} \|x_n - x\|_{H_d^1(\mathbb{T}_\theta^d)} = 0$. From (4.4), we have that the sequence $\{[g(D), 1 \otimes M_{x_n}]\}_{n=0}^\infty$ is Cauchy in the $\mathcal{L}_{d,\infty}$ topology. Hence there is a limit $T \in \mathcal{L}_{d,\infty}$. On the other hand, if $\eta \in C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$ from the Hölder inequality we have:

$$\| (1 \otimes M_{x_n}) \eta - (1 \otimes M_x) \eta \|_{L_2(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N} \leq \| x_n - x \|_{L_d(\mathbb{T}_\theta^d)} \| \eta \|_{L_{2d/(d-2)}(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N}$$

and similarly,

$$\| (1 \otimes M_{x_n}) g(D) \eta - (1 \otimes M_x) g(D) \eta \|_{L_2(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N} \leq \| x_n - x \|_{L_d(\mathbb{T}_\theta^d)} \| g(D) \eta \|_{L_{2d/(d-2)}(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N}.$$

Therefore for each fixed $\eta \in C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$ we have:

$$[g(D), 1 \otimes M_{x_n}] \eta \rightarrow [g(D), 1 \otimes M_x] \eta$$

in the $L_2(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$ sense. Thus,

$$[g(D), 1 \otimes M_x] \eta = T \eta$$

for all $\eta \in C^\infty(\mathbb{T}_\theta^d) \otimes \mathbb{C}^N$. Therefore T and $[g(D), 1 \otimes M_x]$ are equal, and we have:

$$[g(D), 1 \otimes M_{x_n}] \rightarrow [g(D), 1 \otimes M_x]$$

in the $\mathcal{L}_{d,\infty}$ topology. Thus (4.4) holds for all $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. \square

Now we are able to complete the proof of Theorem 1.1.

Proof of Theorem 1.1. Let $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. Combining Lemmas 4.1 and 4.2, we find that

$$\|[\operatorname{sgn}(D), 1 \otimes M_x]\|_{\mathcal{L}_{d,\infty}} \leq C_{d,d}\|x\|_d + B_d\|x\|_{\dot{H}_d^1}.$$

We can remove the dependence on $\|x\|_d$ on the right hand side by the aid of the Poincaré inequality (2.6). Since for constant operator $\hat{x}(0) \in L_\infty(\mathbb{T}_\theta^d)$, it is obvious that $[\operatorname{sgn}(D), 1 \otimes M_{\hat{x}(0)}] = 0$, we have

$$\begin{aligned} \|[\operatorname{sgn}(D), 1 \otimes M_x]\|_{\mathcal{L}_{d,\infty}} &= \|[\operatorname{sgn}(D), 1 \otimes M_{x-\hat{x}(0)}]\|_{\mathcal{L}_{d,\infty}} \\ &\leq C_{d,d}\|x - \hat{x}(0)\|_d + B_d\|x - \hat{x}(0)\|_{\dot{H}_d^1} \\ &\leq C_d\|x\|_{\dot{H}_d^1}. \end{aligned}$$

The theorem is therefore proved. \square

5. PSEUDODIFFERENTIAL OPERATORS ON QUANTUM TORI

In this section we give an introduction to some recent developments in pseudodifferential operators on quantum tori. The most important result stated in this section for us is Theorem 5.6, which is a form of Connes' trace formula obtained in [40].

The theory of pseudodifferential operators goes back to Kohn-Nirenberg [31] and Hörmander [28]. It has been extended to the noncommutative setting, especially the quantum torus case, by many authors; see for instance [39, 32, 34, 23, 57, 59]. Our main references of this part are [6, 1] and [16], while the details can be found in [26, 27]. In the following, let us collect some definitions and well known properties of symbol classes and pseudodifferential operators on quantum tori.

Denote by $\langle \xi \rangle$ the function $(1 + |\xi|^2)^{\frac{1}{2}}$ on \mathbb{R}^d . For every $m \in \mathbb{R}$, the class $S^m(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ consists of all maps $\rho \in C^\infty(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ such that, for all multi-indices $\alpha, \beta \in \mathbb{N}_0^d$, there exists $C_{\alpha,\beta} > 0$ such that

$$\|D^\alpha D_\xi^\beta \rho(\xi)\| \leq C_{\alpha,\beta} \langle \xi \rangle^{m-|\beta|_1}, \quad \forall \xi \in \mathbb{R}^d.$$

Endowed with the locally convex topology generated by the semi-norms

$$p_N^{(m)}(\rho) := \sup_{|\alpha|_1 + |\beta|_1 \leq N} \sup_{\xi \in \mathbb{R}^d} \langle \xi \rangle^{-m+|\beta|_1} \|D^\alpha D_\xi^\beta \rho(\xi)\|, \quad N \in \mathbb{N}_0,$$

$S^m(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ is then a Fréchet space.

Let $\rho \in S^m(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$, $m \in \mathbb{R}$, and $\rho_j(\xi) \in S^{m-j}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ for each $j \in \mathbb{N}$. If for every $N \geq 1$,

$$\rho(\xi) - \sum_{j < N} \rho_j(\xi) \in S^{m-N}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d)),$$

we shall write $\rho(\xi) \sim \sum_{j \geq 0} \rho_j(\xi)$. This is referred to as an asymptotic expansion of the symbol ρ .

The homogeneous class of symbols $\dot{S}^m(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ consists of maps $\rho \in C^\infty(\mathbb{R}^d \setminus \{0\}; C^\infty(\mathbb{T}_\theta^d))$ satisfying

$$\rho(\lambda\xi) = \lambda^m \rho(\xi), \quad \forall \xi \in \mathbb{R}^d \setminus \{0\}, \forall \lambda > 0.$$

In this case, ρ on $\mathbb{R}^d \setminus \{0\}$ is determined by its restriction to \mathbb{S}^{d-1} , the d -dimensional unit sphere. If a (not necessarily homogeneous) symbol ρ admits an asymptotic expansion $\rho \sim \sum_{j \geq 0} \rho_{m-j}$ with $\rho_{m-j} \in \dot{S}^{m-j}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ for each $j \geq 0$, then ρ is called a *classical* symbol, and the leading term ρ_m is called the principal symbol of ρ .

Let us turn to the definition of pseudodifferential operators with the above symbols on quantum torus. Let α_s be a d -parameter group of automorphisms given by

$$(5.1) \quad \alpha_s(U^n) = e^{2\pi i s \cdot n} U^n,$$

which is a periodic version of the action in (2.2) if we identify $[0, 1]^d$ with \mathbb{T}^d by the correspondence $(s_1, s_2, \dots, s_d) \leftrightarrow (e^{2\pi i s_1}, e^{2\pi i s_2}, \dots, e^{2\pi i s_d})$. For $\rho \in C^\infty(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$, let P_ρ be the pseudodifferential operator sending arbitrary $a \in C^\infty(\mathbb{T}_\theta^d)$ to

$$(5.2) \quad P_\rho(a) := \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} e^{-2\pi i s \cdot \xi} \rho(\xi) \alpha_s(a) ds d\xi.$$

Note that this integral does not converge absolutely; it is defined as an oscillatory integral. See [16, 26, 27, 57] for more information. By [26, Proposition 5.9], if $a = \sum_{n \in \mathbb{Z}^d} a_n U^n \in C^\infty(\mathbb{T}_\theta^d)$ and $\rho \in S^m(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ with $m \in \mathbb{R}$, then

$$(5.3) \quad P_\rho(a) = \sum_{n \in \mathbb{Z}^d} \rho(n) a_n U^n,$$

where the sum converges in the operator norm to an element in $C^\infty(\mathbb{T}_\theta^d)$. In other words, the pseudodifferential operator on \mathbb{T}_θ^d with symbol ρ is determined by the value of ρ on \mathbb{Z}^d , which coincides with the definition given in [34].

If $\rho \in S^m(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ with $m \in \mathbb{R}$, then P_ρ is said to be a pseudodifferential operator of order m .

Also note that, by the noncommutativity, if we change the order of $\rho(\xi)$ and $\alpha_s(a)$ in (5.2) (or $\rho(n)$ and U^n in (5.3)), we get another pseudodifferential operator with the same symbol. In [59], these two operators are distinguished as column and row operators. But in this paper, we will not need to consider both kinds of operators, and so we focus only on those with the form (5.2) or (5.3).

Example 5.1. Let us formulate some first examples of symbols defined above.

- i) Let $x \in C^\infty(\mathbb{T}_\theta^d)$ and consider the constant function $\psi(\xi) \equiv x$, $\xi \in \mathbb{R}^d$. Obviously, $\psi \in S^0(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$. So by the above definition, the multiplier $M_x(y) = xy$ on \mathbb{T}_θ^d is an order 0 pseudodifferential operator. The principal symbol of this operator is x itself.
- ii) Let $k \in \mathbb{N}_0^d$. The symbol of the $|k|_1$ -order differential operator $D^k = D_1^{k_1} \cdots D_d^{k_d}$ is $\psi(\xi) = (2\pi\xi_1)^{k_1} (2\pi\xi_2)^{k_2} \cdots (2\pi\xi_d)^{k_d}$. It is easily checked that $\psi \in S^{|k|_1}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$. Thus D^k is a pseudodifferential operator of order $|k|_1$, and its principal symbol is $(2\pi\xi)^k$.
- iii) Let $\alpha \in \mathbb{R}$, and consider the α -order Bessel potential $J^\alpha = (1 - \Delta)^{\frac{\alpha}{2}}$ on the quantum torus, which is a Fourier multiplier with symbol $\psi(\xi) = \langle 2\pi\xi \rangle^\alpha = (1 + |2\pi\xi|^2)^{\frac{\alpha}{2}} \in S^\alpha(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$. Thus, J^α is an α -order pseudodifferential operator. Moreover, as a scalar-valued function, $\psi(\xi)$ has the asymptotic expansion

$$\langle 2\pi\xi \rangle^\alpha \sim \sum_{j=0}^{\infty} \binom{\alpha}{\frac{j}{2}} |2\pi\xi|^{\alpha-2j}.$$

Hence, J^α is classical with principal symbol $|2\pi\xi|^\alpha$. See [26, Proposition 5.14].

The above examples illustrate that both pointwise multipliers M_x and Fourier multipliers T_g from (2.8) are considered as the special cases of pseudodifferential operators. For general symbol ρ , P_ρ may be thought as a limit of linear combinations of operators composed by pointwise multipliers and Fourier multipliers.

The composition of two pseudodifferential operators is again a pseudodifferential operator, and there is a method for computing an asymptotic expansion of its symbol. The following proposition, which is the quantum analogue of the classical result in [56, p. 237], first appears in [16]; a complete proof is given in [27, Proposition 7.5].

Proposition 5.2. *Let ρ_1, ρ_2 be two symbols in $S^{n_1}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ and $S^{n_2}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ respectively. Then there exists a symbol ρ_3 in $S^{n_1+n_2}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ such that*

$$P_{\rho_3} = P_{\rho_1} P_{\rho_2}.$$

Moreover,

$$(5.4) \quad \rho_3 - \sum_{|\alpha|_1 < N_0} \frac{(2\pi i)^{-|\alpha|_1}}{\alpha!} D_\xi^\alpha \rho_1 D^\alpha \rho_2 \in S^{n_1+n_2-N_0}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d)), \quad \forall N_0 \geq 0,$$

where the first derivative D_ξ^α is the derivative of ρ_1 with respect to the variable $\xi \in \mathbb{R}^d$, and the second derivative D^α is the derivation on $C^\infty(\mathbb{T}_\theta^d)$ described in Section 2.2.2.

Many authors have considered the question of the mapping properties of pseudodifferential operators on functions spaces on quantum tori [59, 23, 27]. In this paper we are concerned solely with the boundedness of a pseudodifferential operator on $L_2(\mathbb{T}_\theta^d)$.

The following proposition can be found in [27, Proposition 10.1], [57, Corollary 6.6].

Proposition 5.3. *Let $\rho \in S^0(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$. Then the pseudodifferential operator P_ρ extends to a bounded operator from $L_2(\mathbb{T}_\theta^d)$ to $L_2(\mathbb{T}_\theta^d)$.*

Proposition 5.3 is simply a special case of the general Sobolev space mapping property of pseudodifferential operators [27, Proposition 6.6]. Even greater generalisations to mapping properties of pseudodifferential operators on Sobolev spaces and Besov and Triebel-Lizorkin spaces [59, Section 6.2] are also known.

Symbols of negative order are in particular of order zero, and thus if $m > 0$ and $\rho \in S^{-m}(\mathbb{R}^d, C^\infty(\mathbb{T}_\theta^d))$ then P_ρ has bounded extension on $L_2(\mathbb{T}_\theta^d)$. However in the case of strictly negative order we can provide more detailed information on P_ρ . The following is proved in [27, Lemma 13.6]:

Proposition 5.4. *If $\rho \in S^{-m}(\mathbb{R}^d; C^\infty(\mathbb{T}_\theta^d))$ with $m > 0$, then P_ρ is a compact operator on $L_2(\mathbb{T}_\theta^d)$. Furthermore, $P_\rho \in \mathcal{L}_{\frac{d}{m}, \infty}$.*

The proof of Proposition 5.4 is a simple combination of the fact that since P_ρ has order $-m$, and J^m has order m , the product formula in Proposition 5.2 implies that the composition $P_\rho J^m$ is of order zero. Hence by Proposition 5.3, $P_\rho J^m$ has bounded extension, and since $J^{-m} \in \mathcal{L}_{\frac{d}{m}, \infty}$ (3.3), it follows immediately that $P_\rho \in \mathcal{L}_{\frac{d}{m}, \infty}$.

Thanks to Proposition 5.4, we can easily obtain from the symbol calculus the following:

Corollary 5.5. *Let $x \in C^\infty(\mathbb{T}_\theta^d)$, and $\alpha > 0$. Then*

$$[M_x, (1 - \Delta)^{-\frac{\alpha}{2}}] \in \mathcal{L}_{\frac{d}{\alpha+1}, \infty}.$$

Indeed, $[M_x, (1 - \Delta)^{-\frac{\alpha}{2}}]$ is a pseudodifferential operator of order at most $-\alpha - 1$, as can be seen by a short computation using Proposition 5.2.

Next, we are going to present Connes' trace formula on quantum torus in the specific form obtained in [40, Theorem 6.5]. This trace formula will play a crucial role in the proof of the trace formula for a quantised differential. Recall that if ρ is a homogeneous symbol of order 0, then $\rho(\xi) = \rho(\frac{\xi}{|\xi|})$ for every $\xi \neq 0$. So this ρ could be viewed as a function on the $(d - 1)$ -dimensional sphere \mathbb{S}^{d-1} .

Theorem 5.6. *Let A be a classical pseudodifferential operator on \mathbb{T}_θ^d of order 0 with self-adjoint extension, and denote by ρ_A its principal symbol. Then for any normalised trace φ on $\mathcal{L}_{1, \infty}$, we have*

$$\varphi(|A|^d (1 - \Delta)^{-\frac{d}{2}}) = \frac{1}{d} \int_{\mathbb{S}^{d-1}} \tau(|\rho_A(s)|^d) ds.$$

The reason to refer specifically to [40] is that if d is odd then $|A|^d$ is not a pseudodifferential operator in the usual sense, and so it needs to be understood as an element of the C^* -closure of the algebra of order 0 pseudodifferential operators on $L_2(\mathbb{T}_\theta^d)$. It is proved in [40] that on the C^* -closure the principal symbol mapping extends to a C^* -algebra homomorphism, and hence $\rho_{|A|^d} = |\rho_A|^d$.

6. THE TRACE FORMULA

This section is devoted to the proofs of Theorem 1.2 and Corollary 1.3. That is, we show that for all $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$ and all continuous normalised traces φ on the ideal $\mathcal{L}_{1, \infty}$ that:

$$(6.1) \quad \varphi(|\vec{d}x|^d) = c_d \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} \right) ds$$

for a positive constant c_d . Moreover, there are positive constants $0 < c_d < C_d < \infty$ such that:

$$(6.2) \quad c_d \|x\|_{\dot{H}_d^1}^d \leq \varphi(|\vec{d}x|^d) \leq C_d \|x\|_{\dot{H}_d^1}^d.$$

Our strategy of proof is as follows: first, (6.1) is proved for $x \in C^\infty(\mathbb{T}_\theta^d)$ by aid of the theory of pseudodifferential operators developed in the preceding section. Then by an approximation argument based on the density of $C^\infty(\mathbb{T}_\theta^d)$ in $\dot{H}_d^1(\mathbb{T}_\theta^d)$, we complete the proof of (6.1) in full generality. Finally (6.2) is achieved by bounding the right hand side of (6.1) from above and below by a constant multiple of $\|x\|_{\dot{H}_d^1}^d$.

To begin with, we explain how the operator $|\dot{x}|^d$ can, up to trace class perturbations, be written in the form $|A|^d(1+D^2)^{-\frac{d}{2}}$ for a certain order zero pseudodifferential operator A . Let $x \in C^\infty(\mathbb{T}_\theta^d)$. For $j = 1, \dots, d$, we define the operators $\{A_j\}_{j=1}^d$ on $L_2(\mathbb{T}_\theta^d)$ by

$$A_j \eta := \left(M_{\partial_j x} - \frac{1}{2} \sum_{k=1}^d \left(\frac{D_j D_k}{1-\Delta} M_{\partial_k x} + M_{\partial_k x} \frac{D_j D_k}{1-\Delta} \right) \right) \eta, \quad \eta \in L_2(\mathbb{T}_\theta^d).$$

For each j , A_j is defined initially on $C^\infty(\mathbb{T}_\theta^d)$, but by functional calculus A_j extends uniquely to a bounded operator on $L_2(\mathbb{T}_\theta^d)$ which we denote with the same symbol. We then define the operator A on $\mathbb{C}^N \otimes L_2(\mathbb{T}_\theta^d)$ as

$$(6.3) \quad A := \sum_{j=1}^d \gamma_j \otimes A_j.$$

If $x = x^*$, since ∂_j commutes with the adjoint operation $*$, we have for every $y_1, y_2 \in L_2(\mathbb{T}_\theta^d)$,

$$\langle (\partial_j x) y_1, y_2 \rangle = \tau((\partial_j x) y_1 y_2^*) = \tau(y_1 (\partial_j x^* y_2)^*) = \langle y_1, (\partial_j x^*) y_2 \rangle,$$

which yields $(M_{\partial_j x})^* = M_{\partial_j x^*} = M_{\partial_j x}$. Furthermore, since each D_j is a self-adjoint operator on $L_2(\mathbb{T}_\theta^d)$, we know that

$$\left(\frac{D_j D_k}{1-\Delta} M_{\partial_k x} \right)^* = M_{\partial_k x} \frac{D_j D_k}{1-\Delta}.$$

Therefore, each A_j is a self-adjoint operator, and so is A .

We will now show that $|\dot{x}|^d - |A|^d(1+D^2)^{-\frac{d}{2}} \in \mathcal{L}_1$. The following lemma is an important first step:

Lemma 6.1. *Let $\beta \geq 0$, and $\alpha > 0$ be such that $\alpha + 1 < d$. Then for A defined in (6.3), we have*

$$[|A|, (1+D^2)^{-\frac{\alpha}{2}}] (1+D^2)^{-\frac{\beta}{2}} \in \mathcal{L}_{\frac{d}{\alpha+\beta+1}, \infty}.$$

Proof. As $(1+D^2)^{-\frac{1}{2}} \in \mathcal{L}_{d, \infty}$, the operator $(1+D^2)^{-\frac{\beta}{2}}$ is in $\mathcal{L}_{\frac{d}{\beta}, \infty}$. Thus by Hölder's inequality, it suffices to consider $\beta = 0$. First, we shall prove that $[A_j, (1-\Delta)^{-\frac{\alpha}{2}}] \in \mathcal{L}_{\frac{d}{\alpha+1}, \infty}$. From Corollary 5.5, we have that

$$[M_{\partial_j x}, (1-\Delta)^{-\frac{\alpha}{2}}] \in \mathcal{L}_{\frac{d}{\alpha+1}, \infty}.$$

Hence, by linearity, it suffices to prove that

$$(6.4) \quad \left[\frac{D_j D_k}{1-\Delta} M_{\partial_k x}, (1-\Delta)^{-\frac{\alpha}{2}} \right] \in \mathcal{L}_{\frac{d}{\alpha+1}, \infty}$$

and

$$(6.5) \quad \left[M_{\partial_k x} \frac{D_j D_k}{1-\Delta}, (1-\Delta)^{-\frac{\alpha}{2}} \right] \in \mathcal{L}_{\frac{d}{\alpha+1}, \infty}.$$

Note that (6.5) follows from (6.4) by taking the adjoint. So we prove only (6.4). However, since $\frac{D_j D_k}{1-\Delta}$ commutes with $(1-\Delta)^{-\frac{\alpha}{2}}$, we have

$$\left[\frac{D_j D_k}{1-\Delta} M_{\partial_k x}, (1-\Delta)^{-\frac{\alpha}{2}} \right] = \frac{D_j D_k}{1-\Delta} \left[M_{\partial_k x}, (1-\Delta)^{-\frac{\alpha}{2}} \right].$$

By functional calculus, $\frac{D_j D_k}{1-\Delta}$ is bounded on $L_2(\mathbb{T}_\theta^d)$. Then (6.4) and (6.5) follow from Corollary 5.5 and the boundedness of $\frac{D_j D_k}{1-\Delta}$ on $L_2(\mathbb{T}_\theta^d)$.

Thus, we have proved that

$$[A, (1+D^2)^{-\frac{\alpha}{2}}] = \sum_{j=1}^d \gamma_j \otimes [A_j, (1-\Delta)^{-\frac{\alpha}{2}}] \in \mathcal{L}_{\frac{d}{\alpha+1}, \infty}.$$

To complete the proof, we need to replace A with $|A|$. To this end, we use the result of [47], which implies that if $1 < p < \infty$ and A and B are self-adjoint operators such that $[A, B] \in \mathcal{L}_p$, then $[|A|, B] \in \mathcal{L}_p$; see also [19, Corollary 3.5] for more general results. Since $\frac{d}{\alpha+1} > 1$, the result follows from interpolation. \square

Lemma 6.2. *Let T be a bounded operator on $\mathbb{C}^N \otimes L_2(\mathbb{T}_\theta^d)$, and suppose that*

$$T \in A(1 + D^2)^{-\frac{1}{2}} + \mathcal{L}_{\frac{2d}{3}, \infty},$$

where A is given in (6.3). Then $|T|^d \in \mathcal{L}_{1, \infty}$ and for any continuous normalised trace φ on $\mathcal{L}_{1, \infty}$, we have

$$\varphi(|T|^d) = \varphi(|A|^d(1 + D^2)^{-\frac{d}{2}}).$$

Proof. By the aid of Lemma 6.1, the proof proceeds as in [35, Lemma 14], and is therefore omitted. \square

Lemma 6.3. *For $x \in C^\infty(\mathbb{T}_\theta^d)$ and A defined in (6.3), we have*

$$\dot{d}x - A(1 + D^2)^{-\frac{1}{2}} = [\text{sgn}(D), 1 \otimes M_x] - A(1 + D^2)^{-\frac{1}{2}} \in \mathcal{L}_{\frac{d}{2}, \infty}.$$

Proof. Let $g(D) = D(1 + D^2)^{-\frac{1}{2}}$. Then

$$\begin{aligned} \text{sgn}(D) - g(D) &= \text{sgn}(D) \left(1 - \frac{|D|}{(1 + D^2)^{\frac{1}{2}}} \right) \\ &= \text{sgn}(D) \left(\frac{1}{(1 + D^2)^{\frac{1}{2}} ((1 + D^2)^{\frac{1}{2}} + |D|)} \right). \end{aligned}$$

Since $(1 + D^2)^{-\frac{1}{2}} \in \mathcal{L}_{d, \infty}$, it follows that $\text{sgn}(D) - g(D) \in \mathcal{L}_{\frac{d}{2}, \infty}$. Therefore,

$$[\text{sgn}(D), 1 \otimes M_x] - [g(D), 1 \otimes M_x] \in \mathcal{L}_{\frac{d}{2}, \infty}.$$

Thus, it suffices to prove

$$(6.6) \quad [g(D), 1 \otimes M_x] - A(1 + D^2)^{-\frac{1}{2}} \in \mathcal{L}_{\frac{d}{2}, \infty}.$$

Now let us prove (6.6). By a short computation using Proposition 5.2, we see that the principal symbol of $[\frac{D_j}{(1-\Delta)^{\frac{1}{2}}}, M_x]$ is

$$(6.7) \quad \frac{1}{|2\pi\xi|} \partial_j x - \sum_{k=1}^d \frac{2\pi\xi_k 2\pi\xi_j}{|2\pi\xi|^3} \partial_k x.$$

We also need to determine the principal symbol of $A_j(1 - \Delta)^{-\frac{1}{2}}$, and to this end we compute the principal symbol of A_j . Recall that

$$A_j = M_{\partial_j x} - \frac{1}{2} \sum_{k=1}^d \left(\frac{D_j D_k}{1 - \Delta} M_{\partial_k x} + M_{\partial_k x} \frac{D_j D_k}{1 - \Delta} \right).$$

It is evident that the symbol of $M_{\partial_k x} \frac{D_j D_k}{1 - \Delta}$ is $\frac{2\pi\xi_j 2\pi\xi_k}{1 + |2\pi\xi|^2} \partial_k x$, so the principal symbol is $\frac{\xi_j \xi_k}{|\xi|^2} \partial_k x$. By Proposition 5.2, we know that the symbol of $\frac{D_j D_k}{1 - \Delta} M_{\partial_k x}$ has the asymptotic expansion

$$\sum_{\alpha \in \mathbb{N}_0^d} \frac{(2\pi i)^{-|\alpha|_1}}{\alpha!} D_\xi^\alpha \left(\frac{2\pi\xi_j 2\pi\xi_k}{(1 + |2\pi\xi|^2)^{\frac{1}{2}}} \right) D^\alpha(\partial_k x).$$

Thus, the principal symbol of $\frac{1}{2} \sum_{k=1}^d \left(\frac{D_j D_k}{1 - \Delta} M_{\partial_k x} + M_{\partial_k x} \frac{D_j D_k}{1 - \Delta} \right)$ is $\sum_{k=1}^d \frac{\xi_k \xi_j}{|\xi|^2} \partial_k x$, which ensures that the principal symbol of $A_j(1 - \Delta)^{-\frac{1}{2}}$ is of order -1 , given by

$$\frac{1}{2\pi|\xi|} \partial_j x - \sum_{k=1}^d \frac{\xi_k \xi_j}{2\pi|\xi|^3} \partial_k x,$$

the same as that of $[\frac{D_j}{(1-\Delta)^{\frac{1}{2}}}, M_x]$ given in (6.7). Hence, the order of $[\frac{D_j}{(1-\Delta)^{\frac{1}{2}}}, M_x] - A_j(1-\Delta)^{-\frac{1}{2}}$ is -2 . By Theorem 5.4, we have

$$[\frac{D_j}{(1-\Delta)^{\frac{1}{2}}}, M_x] - A_j(1-\Delta)^{-\frac{1}{2}} \in \mathcal{L}_{\frac{d}{2}, \infty}.$$

Since $[g(D), 1 \otimes M_x] = \sum_j \gamma_j \otimes [\frac{D_j}{(1-\Delta)^{\frac{1}{2}}}, M_x]$ and $A(1+D^2)^{-\frac{1}{2}} = \sum_j \gamma_j \otimes A_j(1-\Delta)^{-\frac{1}{2}}$, we obtain (6.6). The lemma is thus proved. \square

Based on the above lemmas, we are able to complete the proof of Theorem 1.2.

Proof of Theorem 1.2. Assume initially that $x \in C^\infty(\mathbb{T}_\theta^d)$. By Lemma 6.3, we have that:

$$\bar{d}x \in A(1+D^2)^{-\frac{1}{2}} + \mathcal{L}_{\frac{d}{2}, \infty}.$$

For any continuous normalised trace φ on $\mathcal{L}_{1, \infty}$, we invoke Lemma 6.2 to obtain that:

$$\varphi(|\bar{d}x|^d) = \varphi(|A|^d(1+D^2)^{-\frac{d}{2}}).$$

In the proof of Lemma 6.3, we have that the principal symbol of A_j is $\partial_j x - \sum_{k=1}^d \frac{\xi_k \xi_j}{|\xi|^2} \partial_k x$, which restricted to the unit sphere \mathbb{S}^{d-1} is $\partial_j x - \sum_{k=1}^d \xi_k \xi_j \partial_k x$. Now we appeal to Theorem 5.6 to conclude

$$\varphi(|\bar{d}x|^d) = c_d \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} \right) ds.$$

However the appeal to Theorem 5.6 relies on the assumption that $x \in C^\infty(\mathbb{T}_\theta^d)$, so we remove this assumption by an approximation argument. Indeed, let $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. By Theorem 1.1, we have $\bar{d}x \in \mathcal{L}_{d, \infty}$. By the density of $C^\infty(\mathbb{T}_\theta^d)$ in $\dot{H}_d^1(\mathbb{T}_\theta^d)$ (see [60, Proposition 2.7]), we can choose a sequence $\{x_n\}_{n=1}^\infty \subset C^\infty(\mathbb{T}_\theta^d)$ such that $x_n \rightarrow x$ in $\dot{H}_d^1(\mathbb{T}_\theta^d)$. We shall show that $\varphi(|\bar{d}x_n|^d) \rightarrow \varphi(|\bar{d}x|^d)$ and

$$(6.8) \quad \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x_n - s_j \sum_{k=1}^d s_k \partial_k x_n|^2 \right)^{\frac{d}{2}} \right) ds \rightarrow \int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{d}{2}} \right) ds.$$

Note that we have a bound:

$$\int_{\mathbb{S}^{d-1}} \tau \left(\left(\sum_{j=1}^d |\partial_j x_n - s_j \sum_{k=1}^d s_k \partial_k x_n|^2 \right)^{\frac{d}{2}} \right) ds \leq C_d \|x_n\|_{\dot{H}_d^1}$$

for a certain constant C_d , and hence (6.8) is immediate. On the other hand, using Theorem 1.1, we have:

$$\|\bar{d}x - \bar{d}x_n\|_{\mathcal{L}_{d, \infty}} \leq C_d \|x - x_n\|_{\dot{H}_d^1(\mathbb{T}_\theta^d)} \rightarrow 0.$$

By a verbatim repetition of the argument in the proof of [35, Theorem 17], we get

$$\| |\bar{d}x|^d - |\bar{d}x_n|^d \|_{\mathcal{L}_{1, \infty}} \rightarrow 0.$$

Since the trace φ is assumed to be continuous in the $\mathcal{L}_{1, \infty}$ quasi-norm, it follows that $\varphi(|\bar{d}x_n|^d) \rightarrow \varphi(|\bar{d}x|^d)$. \square

We are now concerned with relating the right hand side of the trace formula in Theorem 1.2 with the \dot{H}_d^1 -norm of x .

Proof of Corollary 1.3. We prove the upper bound first. Denote

$$T(x) := \left(\sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2 \right)^{\frac{1}{2}}, \quad s \in \mathbb{S}^{d-1}.$$

Then

$$\begin{aligned}
|T(x)|^2 &= \sum_{j=1}^d \left| \partial_j x - s_j \sum_{k=1}^d s_k \partial_k x \right|^2 \\
&= \sum_{j=1}^d \left(|\partial_j x|^2 - \sum_{k=1}^d (s_j \partial_j x^* \cdot s_k \partial_k x + s_k \partial_k x^* \cdot s_j \partial_j x) + s_j^2 \left| \sum_{k=1}^d s_k \partial_k x \right|^2 \right) \\
&= \sum_{j=1}^d |\partial_j x|^2 + \left| \sum_{k=1}^d s_k \partial_k x \right|^2 - \sum_{j,k=1}^d (s_j \partial_j x^* \cdot s_k \partial_k x + s_k \partial_k x^* \cdot s_j \partial_j x).
\end{aligned}$$

However, observing that

$$\left| \sum_{j=1}^d s_j \partial_j x \right|^2 = \sum_{j,k=1}^d s_j \partial_j x^* \cdot s_k \partial_k x,$$

we get

$$(6.9) \quad |T(x)|^2 = \sum_{j=1}^d |\partial_j x|^2 - \left| \sum_{j=1}^d s_j \partial_j x \right|^2.$$

We then have have easily:

$$|T(x)|^2 \leq \sum_{j=1}^d |\partial_j x|^2.$$

Therefore,

$$\| |T(x)|^2 \|_{\frac{d}{2}} \leq \left\| \sum_{j=1}^d |\partial_j x|^2 \right\|_{\frac{d}{2}}.$$

Hence, by (2.5), for every $s \in \mathbb{S}^{d-1}$, we have

$$\tau(|T(x)|^d) = \| |T(x)|^2 \|_{\frac{d}{2}}^{\frac{d}{2}} \leq \left\| \sum_{j=1}^d |\partial_j x|^2 \right\|_{\frac{d}{2}}^{\frac{d}{2}} \leq C_d \|x\|_{H_d^1}^d.$$

Thus the upper bound is proved.

Now we prove the lower bound. Since $|T(x)|^2 = \sum_{j=1}^d |\partial_j x - s_j \sum_{k=1}^d s_k \partial_k x|^2$, for each j we have

$$\left| \partial_j x - s_j \sum_{k=1}^d s_k \partial_k x \right|^2 \leq |T(x)|^2,$$

and therefore,

$$(6.10) \quad \left\| \partial_j x - s_j \sum_{k=1}^d s_k \partial_k x \right\|_d \leq \|T(x)\|_d.$$

For brevity, define

$$X_j = \left\| \partial_j x - s_j \sum_{k=1}^d s_k \partial_k x \right\|_d.$$

Then (6.11) implies that

$$(6.11) \quad \left(\sum_{j=1}^d X_j \right)^d \leq d^d \|T(x)\|_d^d.$$

By the triangle inequality,

$$\begin{aligned}
X_j &= \left\| (1 - s_j^2) \partial_j x - \sum_{k \neq j} s_j s_k \partial_k x \right\|_d \\
&\geq (1 - s_j^2) \|\partial_j x\|_d - \sum_{k \neq j} |s_j s_k| \|\partial_k x\|_d.
\end{aligned}$$

and therefore,

$$\sum_{j=1}^d X_j \geq \sum_{j=1}^d \left((1 - s_j^2) - \sum_{k \neq j} |s_j s_k| \right) \|\partial_j x\|_d.$$

Now, select $1 \leq l \leq d$ such that $\|\partial_l x\|_d$ is the minimum of $\{\|\partial_1 x\|_d, \|\partial_2 x\|_d, \dots, \|\partial_d x\|_d\}$. Denote by e_l the l -th canonical basic vector of \mathbb{R}^d , and assume that $s \in B(e_l, \varepsilon) \cap \mathbb{S}^{d-1}$. We have:

$$\left| (1 - s_l^2) - \sum_{k \neq l} |s_l s_k| \right| \leq \max \left(1 - s_l^2, \sum_{k \neq l} |s_k s_l| \right)$$

Hence, $(1 - s_l) \leq |s - e_l| \leq \varepsilon$, so $(1 - s_l^2) = (1 - s_l)(1 + s_l) \leq 2\varepsilon$, and by the Cauchy-Schwarz inequality

$$\begin{aligned} \sum_{k \neq l} |s_k s_l| &\leq \left(\sum_{k \neq l} |s_k|^2 \right)^{\frac{1}{2}} d^{\frac{1}{2}} |s_l| \\ &\leq |s - e_l| d^{\frac{1}{2}} |s_l| \\ &\leq \sqrt{d} \varepsilon. \end{aligned}$$

So,

$$(6.12) \quad \left| (1 - s_l^2) - \sum_{k \neq l} |s_l s_k| \right| \leq \max\{2, \sqrt{d}\} \varepsilon.$$

On the other hand, if $j \neq l$, then $|s_j| \leq \varepsilon$ and so:

$$(6.13) \quad \begin{aligned} (1 - s_j^2) - \sum_{k \neq j} |s_k s_j| &= 1 - |s_j| \sum_{k=1}^d |s_k| \\ &\geq 1 - \sqrt{d} \varepsilon. \end{aligned}$$

If we select ε sufficiently small, we have $1 - \sqrt{d} \varepsilon \geq 3 \max\{2, \sqrt{d}\} \varepsilon$. Then combining (6.12) and (6.13), we have that for all $j \neq l$:

$$3 \left| (1 - s_l^2) - \sum_{k \neq l} |s_l s_k| \right| \leq (1 - s_j^2) - \sum_{k \neq j} |s_k s_j|,$$

and thus,

$$\left| (1 - s_l^2) - \sum_{k \neq l} |s_l s_k| \right| \|\partial_l x\|_d \leq \frac{1}{3} \left((1 - s_j^2) - \sum_{k \neq j} |s_k s_j| \right) \|\partial_j x\|_d.$$

Therefore, using the numerical inequality that if $|z| \leq \frac{1}{3}|w|$ then $|z - w| \geq \frac{2}{3}|w|$, we have

$$\begin{aligned} &\left((1 - s_j^2) - \sum_{k \neq j} |s_k s_j| \right) \|\partial_j x\|_d + \left((1 - s_l^2) - \sum_{k \neq l} |s_l s_k| \right) \|\partial_l x\|_d \\ &\geq \frac{2}{3} \left((1 - s_j^2) - \sum_{k \neq j} |s_k s_j| \right) \|\partial_j x\|_d \\ &\geq \frac{1}{3} (1 - \sqrt{d} \varepsilon) (\|\partial_j x\|_d + \|\partial_l x\|_d). \end{aligned}$$

Consequently, for $s \in B(e_l, \varepsilon) \cap \mathbb{S}^{d-1}$, we have,

$$\sum_{j=1}^d X_j \geq (1 - \sqrt{d} \varepsilon) \sum_{j=1}^d \|\partial_j x\|_d.$$

Now,

$$\begin{aligned} \int_{\mathbb{S}^{d-1}} \left(\sum_{j=1}^d X_j \right)^d ds &\geq \int_{B(e_l, \varepsilon) \cap \mathbb{S}^{d-1}} \left(\sum_{j=1}^d X_j \right)^d ds \\ &\geq c_{d, \varepsilon} \|x\|_{\dot{H}_d^1}. \end{aligned}$$

By virtue of (6.11), the desired conclusion is proved. \square

7. PROOF OF THEOREM 1.4

In this section, we are going to give the proof of Theorem 1.4. We require a lemma on the quantised derivative of x acting by a Fourier multiplier.

Recall that for a function $\psi \in L_1(\mathbb{T}^d)$, the convolution with $x \in L_2(\mathbb{T}_\theta^d)$ is defined as:

$$\psi * x = \int_{\mathbb{T}^d} \psi(w) \alpha_{w^{-1}}(x) dm(w).$$

Lemma 7.1. *Let $\psi \in L_1(\mathbb{T}^d)$. If $x \in L_2(\mathbb{T}_\theta^d)$ is such that $\check{d}x$ extends to a bounded operator in $\mathcal{L}_{d,\infty}$, then $\check{d}(\psi * x)$ also extends to a bounded operator in $\mathcal{L}_{d,\infty}$ and we have:*

$$\|\check{d}(\psi * x)\|_{\mathcal{L}_{d,\infty}} \leq C_d \|\check{d}x\|_{\mathcal{L}_{d,\infty}} \|\psi\|_1$$

for a certain constant C_d .

Proof. Let α be the d -parameter group of automorphisms given in (2.2), i.e. if $u \in \mathbb{T}^d$ then $\alpha_u(U^n) = u^n U^n$. Then for each $u \in \mathbb{T}^d$, α_u commutes with Fourier multipliers on \mathbb{T}_θ^d , and $u \mapsto \alpha_u$ is a strongly continuous family of unitary operators on $L_2(\mathbb{T}_\theta^d)$. By the definition of convolution, we have

$$(7.1) \quad \psi * x = \int_{\mathbb{T}^d} \psi(u) \alpha_u^{-1}(x) dm(u).$$

Since α_u and $\frac{D_j}{\sqrt{D_1^2 + D_2^2 + \dots + D_d^2}}$ commute, we see that $1 \otimes \alpha_u$ commutes with $\text{sgn}(D)$. Therefore, by the fact that $(\alpha_u^{-1}(x))y = \alpha_u^{-1}(x(\alpha_u(y)))$, we obtain

$$[\text{sgn}(D), 1 \otimes M_{\psi * x}] = \int_{\mathbb{T}^d} \psi(u) (1 \otimes \alpha_{u^{-1}})[\text{sgn}(D), 1 \otimes M_x](1 \otimes \alpha_u) dm(u).$$

Applying [35, Lemma 18] to the finite Borel measure $\psi(u) dm(u)$ on \mathbb{T}^d , we get

$$\|[\text{sgn}(D), 1 \otimes M_{\psi * x}]\|_{\mathcal{L}_{d,\infty}} \leq C_d \|[\text{sgn}(D), 1 \otimes M_x]\|_{\mathcal{L}_{d,\infty}} \|\psi\|_1$$

where the constant comes from the use of the quasi-triangle inequality in the $\mathcal{L}_{d,\infty}$ quasi-norm. This now completes the proof. \square

Proof of Theorem 1.4. Firstly, we prove the theorem for self-adjoint $x \in L_2(\mathbb{T}_\theta^d)$. If we show that $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$, then Corollary 1.3 will ensure that there exists a constant $c_d > 0$ such that for all continuous normalised traces φ on $\mathcal{L}_{1,\infty}$,

$$c_d \|x\|_{\dot{H}_d^1} \leq \varphi(|\check{d}x|^d)^{\frac{1}{d}} \leq \|\check{d}x\|_{d,\infty}.$$

Thus, we are reduced to proving $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$.

Consider the square Fejér mean

$$F_N(x) = \sum_{m \in \mathbb{Z}^d, \max_j |m_j| \leq N} \left(1 - \frac{|m_1|}{N+1}\right) \dots \left(1 - \frac{|m_d|}{N+1}\right) \hat{x}(m) U^m.$$

For every $N \in \mathbb{N}$, it is the convolution of x with the periodic function

$$F_N(u) = \frac{1}{(N+1)^d} \left(\frac{\sin(\pi(N+1)u_1)}{\sin(\pi u_1)}\right)^2 \dots \left(\frac{\sin(\pi(N+1)u_d)}{\sin(\pi u_d)}\right)^2.$$

The family $\{F_N\}_{N \in \mathbb{N}}$ is an approximate identity of $L_1(\mathbb{T}^d)$ (see [25]), so we have uniform bound of $\|F_N\|_1$ in $N \in \mathbb{N}$. Thus we can apply Lemma 7.1 to F_N . The result is

$$\|\check{d}(F_N(x))\|_{\mathcal{L}_{d,\infty}} \leq \|\check{d}x\|_{\mathcal{L}_{d,\infty}} \|F_N\|_1 \leq C \|\check{d}x\|_{\mathcal{L}_{d,\infty}}.$$

Since each $F_N(x)$ is a polynomial in \mathbb{T}_θ^d , Corollary 1.3 yields

$$c_d \|F_N(x)\|_{\dot{H}_d^1} \leq \varphi(|\check{d}(F_N(x))|^d)^{\frac{1}{d}} \leq C \|\check{d}x\|_{\mathcal{L}_{d,\infty}}.$$

Hence, for each j , we obtain a bounded sequence $\{\partial_j F_N(x)\}_{N \in \mathbb{N}}$ in $L_d(\mathbb{T}_\theta^d)$. Moreover, since $L_d(\mathbb{T}_\theta^d)$ is reflexive, we may assume that $\partial_j F_N(x)$ converges to some $y_j \in L_d(\mathbb{T}_\theta^d)$. On the other hand, by [5, Proposition 3.1], we have $\lim_N F_N(x) = x$ in $L_2(\mathbb{T}_\theta^d)$. Hence, $\partial_j F_N(x)$ converges to $\partial_j x$ in $\mathcal{D}'(\mathbb{T}_\theta^d)$. Therefore, we have $y_j = \partial_j x \in L_d(\mathbb{T}_\theta^d)$. Consequently, we conclude that $x \in \dot{H}_d^1(\mathbb{T}_\theta^d)$.

It remains to consider $x \in L_2(\mathbb{T}_\theta^d)$ which are not self-adjoint. Write $x = x_1 + ix_2$ with

$$x_1 = \frac{1}{2}(x + x^*), \quad x_2 = \frac{1}{2i}(x - x^*).$$

If $[\text{sgn}(D), 1 \otimes M_x] \in \mathcal{L}_{d,\infty}$, then $[\text{sgn}(D), 1 \otimes M_{x^*}] = -[\text{sgn}(D), 1 \otimes M_x]^* \in \mathcal{L}_{d,\infty}$. Then we have $[\text{sgn}(D), 1 \otimes M_{x_1}] \in \mathcal{L}_{d,\infty}$ and $[\text{sgn}(D), 1 \otimes M_{x_2}] \in \mathcal{L}_{d,\infty}$. By the above conclusion for self-adjoint elements, we know that $x_1, x_2 \in \dot{H}_d^1(\mathbb{T}_\theta^d)$, which implies $x = x_1 + ix_2 \in \dot{H}_d^1(\mathbb{T}_\theta^d)$. More precisely,

$$\begin{aligned} \|x\|_{\dot{H}_d^1} &\leq \|x_1\|_{\dot{H}_d^1} + \|x_2\|_{\dot{H}_d^1} \\ &\leq C_1(\|[\text{sgn}(D), 1 \otimes M_{x_1}]\|_{\mathcal{L}_{d,\infty}} + \|[\text{sgn}(D), 1 \otimes M_{x_2}]\|_{\mathcal{L}_{d,\infty}}) \\ &\leq C_2(\|[\text{sgn}(D), 1 \otimes M_x]\|_{\mathcal{L}_{d,\infty}} + \|[\text{sgn}(D), 1 \otimes M_x]^*\|_{\mathcal{L}_{d,\infty}}) \\ &= 2C_2\|[\text{sgn}(D), 1 \otimes M_x]\|_{\mathcal{L}_{d,\infty}}. \end{aligned}$$

The theorem is proved. \square

Acknowledgements. The authors wish to thank the anonymous referees for careful reading and useful suggestions; in particular one referee pointed out how our main results could be proved without an L_∞ condition. We are also greatly indebted to Professor Raphaël Ponge for many helpful comments on the section of Pseudodifferential Operators. The authors are supported by Australian Research Council (grant no. FL170100052); X. Xiong is also partially supported by the National Natural Science Foundation of China (grant no. 11301401).

REFERENCES

- [1] S. Baaj. Calcul pseudodifférentiel et produits croisés de C^* -algèbres, I and II. *C. R. Acad. Sc. Paris, sér. I*, 307 (1988), 581–586 and 663–666.
- [2] J. Bellissard. K -theory of C^* -algebras in solid state physics. In *Statistical mechanics and field theory: mathematical aspects (Groningen, 1985)*, volume 257 of *Lecture Notes in Phys.*, pages 99–156. Springer, Berlin, 1986.
- [3] J. Bellissard, A. van Elst, and H. Schulz-Baldes. The noncommutative geometry of the quantum Hall effect. *J. Math. Phys.*, 35(10):5373–5451, 1994. Topology and physics.
- [4] M. S. Birman, M. Z. Solomyak. Operator integration, perturbations and commutators. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 170* (1989) 34–66, 321.
- [5] Z. Chen, Q. Xu, and Z. Yin. Harmonic analysis on quantum tori. *Comm. Math. Phys.*, 322 (2013), 755805.
- [6] A. Connes. C^* -algèbres et géométrie différentielle. *C. R. Acad. Sc. Paris, sér. A*, 290 (1980), 599–604.
- [7] A. Connes. Noncommutative differential geometry. *Inst. Hautes Études Sci. Publ. Math.*, (62):257–360, 1985.
- [8] A. Connes. The action functional in noncommutative geometry. *Comm. Math. Phys.*, 117 (1988), 673–683.
- [9] A. Connes. *Noncommutative Geometry*. Academic Press, San Diego, CA, 1994.
- [10] A. Connes. Noncommutative geometry and reality. *J. Math. Phys.*, 36 (1995), no. 11 6194–6231.
- [11] A. Connes, M. R. Douglas, and A. Schwarz. Noncommutative geometry and matrix theory: compactification on tori. *J. High Energy Phys.*, (2):Paper 3, 35, 1998.
- [12] A. Connes, E. McDonald, F. Sukochev and D. Zanin Conformal trace theorem for Julia sets of quadratic polynomials. *Ergodic Theory and Dynamical Systems*, 1–26. (2017)
- [13] A. Connes and H. Moscovici. Modular curvature for noncommutative two-tori. *J. Amer. Math. Soc.*, 27 (2014), 639–684.
- [14] A. Connes, F. Sukochev and D. Zanin. Trace theorem for quasi-Fuchsian groups. *Mat. Sb.*, 208 (2017), 59–90.
- [15] A. Connes, D. Sullivan and N. Teleman. Quasiconformal mappings, operators on Hilbert space, and local formulae for characteristic classes. *Topology*, 33 (1994), 663–681.
- [16] A. Connes and P. Tretkoff. The Gauss-Bonnet theorem for the noncommutative two torus. *Noncommutative geometry, arithmetic, and related topics*, Johns Hopkins Univ. Press, Baltimore, MD, 2011, 141158.
- [17] M. Cwikel. Weak type estimates for singular values and the number of bound states of Schrödinger operators. *Ann. of Math.*, 106 (1977), 93–100
- [18] P.G. Dodds, T.K. Dodds and B. de Pagter. Fully symmetric operator spaces. *Integral Equations Operator Theory*, 15 (1992), 942–972.
- [19] P.G. Dodds, T.K. Dodds, B. de Pagter and F. Sukochev. Lipschitz continuity of the absolute value and Riesz projections in symmetric operator spaces. *J. Funct. Anal.*, 148 (1997), 28–69.
- [20] K. Dykema, T. Figiel, G. Weiss and M. Wodzicki. Commutator structure of operator ideals. *Adv. Math.* 185 (2004), 1–79.
- [21] E. G. Effros and F. Hahn. Locally compact transformation groups and C^* -algebras. *Bull. Amer. Math. Soc.* 73 (1967), 222–226.
- [22] L. Gheorghie. Hankel operators in Schatten ideals. *Ann. Mat. Pura Appl.* (4) 180 (2001), no. 2, 203–210.
- [23] A. M. González-Pérez, M. Junge and J. Parcet. Singular integral in quantum Euclidean spaces. arXiv:1705.01081.

- [24] J. M. Gracia-Bonda, J. Vrilly and H. Figueroa Elements of noncommutative geometry. Birkhuser Advanced Texts: Basler Lehrbcher. [Birkhuser Advanced Texts: Basel Textbooks] Birkhuser Boston, Inc., Boston, MA, 2001. xviii+685 pp.
- [25] L. Grafakos. *Classical Fourier analysis, Second Edition*. Springer-Verlag, New York, 2008.
- [26] H. Ha, G. Lee and R. Ponge. Pseudodifferential calculus on noncommutative tori, I. *Internat. J. Math.* 30 (2019), no. 8, 1950033, 74 pp.
- [27] H. Ha, G. Lee and R. Ponge. Pseudodifferential calculus on noncommutative tori, II. *Internat. J. Math.* 30 (2019), no. 8, 1950034, 73 pp.
- [28] L. Hörmander. Pseudo-differential operators. Vol. 18 (1965), Issue 3, 501-517.
- [29] S. Janson and T.H. Wolff. Schatten classes and commutators of singular integral operators. *Ark. Mat.*, 20 (1982), 301–310.
- [30] N. Kalton, Nigel, S. Lord, D. Potapov and F. Sukochev. Traces of compact operators and the noncommutative residue. *Adv. Math.* 235 (2013), 1–55.
- [31] J. J. Kohn and L. Nirenberg. An algebra of pseudo-differential operators. *Comm. Pure Appl. Math.* 18 (1965), 269-305.
- [32] M. Lein, M. Măntoiu and S. Richard. Magnetic pseudodifferential operators with coefficients in C^* -algebras. *Publ. Res. Inst. Math. Sci.*, 46 (2010), 755-788.
- [33] G. Levitina, F. Sukochev and D. Zanin. Cwikel estimates revisited. arXiv:1703.04254.
- [34] C. Lévy, C.N. Jiménez and S. Paycha. The canonical trace and the noncommutative residue on the noncommutative torus. *Trans. Amer. Math. Soc.* 368 (2016), no. 2, 1051-1095.
- [35] S. Lord, E. McDonald, F. Sukochev and D. Zanin. Quantum differentiability of essentially bounded functions on Euclidean space. *J. Funct. Anal.*, 273 (2017), 2353-2387.
- [36] S. Lord, D. Potapov and F. Sukochev. Measures from Dixmier traces and zeta functions. *J. Funct. Anal.*, 259 (2010), 1915–1949.
- [37] S. Lord, F. Sukochev and D. Zanin. *Singular traces: theory and applications*. Walter de Gruyter, Vol. 46, 2012.
- [38] S. Lord, F. Sukochev and D. Zanin. Advances in Dixmier traces and applications. (submitted manuscript).
- [39] M. Măntoiu, R. Purice and S. Richard. Twisted crossed products and magnetic pseudodifferential operators, in Advances in operator algebras and mathematical physics. *Theta Ser. Adv. Math.* 5 (2005), 137-172.
- [40] E. McDonald, F. Sukochev and D. Zanin. A C^* -algebraic approach to the principal symbol II. *Math. Ann.* 374 (2019), no. 1-2, 273-322.
- [41] B. de Pagter, F. Sukochev and H. Witvliet. Double operator integrals. *J. Funct. Anal.*, 192 (2002), 52111.
- [42] V. V. Peller. *Hankel Operators and Their Applications*. Springer Monographs in Mathematics, 2003.
- [43] M. Pimsner and D. Voiculescu. Exact sequences for K -groups and Ext-groups of certain cross-product C^* -algebras. *J. Operator Theory*, 4(1):93–118, 1980.
- [44] G. Pisier and Q. Xu. Noncommutative L^p -spaces. *Handbook of the geometry of Banach spaces Vol. 2*, ed. W. B. Johnson and J. Lindenstrauss, 2003, 1459-1517, North-Holland, Amsterdam.
- [45] D. Potapov and F. Sukochev. Lipschitz and commutator estimates in symmetric operator spaces. *J. Operator Theory*, 59:1(2008), 211234.
- [46] D. Potapov and F. Sukochev. Unbounded Fredholm modules and double operator integrals. *J. Reine Angew. Math.*, 626 (2009) 159185.
- [47] D. Potapov and F. Sukochev. Operator-Lipschitz functions in Schatten-von Neumann classes. *Acta Math.*, 207 (2011), 375-389.
- [48] M. A. Rieffel. C^* -algebras associated with irrational rotations. *Pacific J. Math.*, 93(2):415–429, 1981.
- [49] M A. Rieffel. Deformation quantization for actions of \mathbf{R}^d . *Mem. Amer. Math. Soc.*, 106(506):x+93, 1993.
- [50] É. Ricard and Q. Xu. Complex interpolation of weighted noncommutative L_p -spaces. *Houston J. Math.* , 37 (2011),1165–1179.
- [51] R. Rochberg and S. Semmes. Nearly weakly orthonormal sequences, singular value estimates, and Calderón–Zygmund operators. *J. Funct. Anal.*, 86 (1989), 237-306.
- [52] N. Seiberg and E. Witten. String theory and noncommutative geometry. *J. High Energy Phys.*, (9):Paper 32, 93, 1999.
- [53] E. Semenov, F. Sukochev, A. Usachev and D. Zanin Banach limits and traces on $\mathcal{L}_{1,\infty}$. *Adv. Math.* 285 (2015), 568–628.
- [54] B. Simon. *Trace Ideals and Their Applications*. London Mathematical Society Lecture Note Series, vol.35, Cambridge University Press, CambridgeNew York, 1979.
- [55] M. Spera. Sobolev theory for noncommutative tori. *Rend. Sem. Mat. Univ. Padova.* 86 (1992), 143-156.
- [56] E. M. Stein. Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [57] J. Tao. The theory of pseudo-differential operators on the noncommutative n -torus. *2018 J. Phys.: Conf. Ser.* 965(2018) 1-12.
- [58] J. Xia. Geometric invariants of the quantum Hall effect. *Comm. Math. Phys.*, 119(1):29–50, 1988.
- [59] R. Xia and X. Xiong. Mapping properties of operator-valued pseudo-differential operators. arXiv:1804.03435.
- [60] X. Xiong, Q. Xu and Z. Yin. Sobolev, Besov and Triebel-Lizorkin spaces on quantum tori. *Mem. Amer. Math. Soc.* 252 (2018), no. 1203, vi+118 pp..
- [61] Q. Xu. Noncommutative L_p -spaces and martingale inequalities. (unpublished manuscript), 2007.

SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, KENSINGTON, NSW 2052, AUSTRALIA
E-mail address: `edward.mcdonald@unsw.edu.au`

SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, KENSINGTON, NSW 2052, AUSTRALIA
E-mail address: `f.sukochev@unsw.edu.au`

INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, 150001 HARBIN, CHINA,
AND SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, KENSINGTON, NSW 2052, AUSTRALIA
E-mail address: `xxiong@hit.edu.cn`